

EGERVÁRY RESEARCH GROUP
ON COMBINATORIAL OPTIMIZATION



TECHNICAL REPORTS

TR-2010-13. Published by the Egerváry Research Group, Pázmány P. sétány
1/C, H-1117, Budapest, Hungary. Web site: www.cs.elte.hu/egres. ISSN
1587-4451.

Tree metrics and edge-disjoint S -paths

Hiroshi Hirai and Gyula Pap

December, 2010 (revised October, 2012)

Tree metrics and edge-disjoint S -paths

Hiroshi Hirai ^{*} and Gyula Pap ^{**}

Abstract

Given an undirected graph $G = (V, E)$ with a terminal set $S \subseteq V$, a weight function $\mu : \binom{S}{2} \rightarrow \mathbf{Z}_+$ on terminal pairs, and an edge-cost $a : E \rightarrow \mathbf{Z}_+$, the μ -weighted minimum-cost edge-disjoint S -paths problem (μ -CEDP) is to maximize $\sum_{P \in \mathcal{P}} \mu(s_P, t_P) - a(P)$ over all edge-disjoint sets \mathcal{P} of S -paths, where s_P, t_P denote the ends of P and $a(P)$ is the sum of edge-cost $a(e)$ over edges e in P .

Our main result is a complete characterization of terminal weights μ for which μ -CEDP is tractable and admits a combinatorial min-max theorem. We prove that if μ is a tree metric, then μ -CEDP is solvable in polynomial time and has a combinatorial min-max formula, which extends Mader's edge-disjoint S -paths theorem and its minimum-cost generalization by Karzanov. Our min-max theorem includes the dual half-integrality, which was earlier conjectured by Karzanov for a special case. We also prove that μ -EDP, which is μ -CEDP with $a = 0$, is NP-hard if μ is not a truncated tree metric, where a truncated tree metric is a weight function represented as pairwise distances between balls in a tree. On the other hand, μ -CEDP for a truncated tree metric μ reduces to μ' -CEDP for a tree metric μ' . Thus our result is best possible unless $P = NP$. As an application, we get a good approximation algorithm for μ -EDP with "near" tree metric μ by utilizing results from the theory of low-distortion embedding.

1 Introduction

Let G be an undirected graph with vertex set $V = V(G)$ and edge set $E = E(G)$. We are given a specified set S of vertices, called *terminals*. An S -path is a path connecting distinct terminals in S . Let $a : E \rightarrow \mathbf{Z}_+$ be a nonnegative integral edge-cost, and let $\mu : \binom{S}{2} \rightarrow \mathbf{Z}_+$ be a nonnegative integral terminal weight defined on the set $\binom{S}{2}$ of all (unordered) pairs of S . One (s, t) -path P has the value $\mu(s, t)$ as well as the cost $a(P) := \sum_{e \in E} a(e)$. We want to find an edge-disjoint set \mathcal{P} of

^{*}Department of Mathematical Informatics, University of Tokyo, Tokyo 113-8656, Japan, hirai@mist.i.u-tokyo.ac.jp

^{**}MTA-ELTE Egerváry Research Group (EGRES). Department of Operations Research, Eötvös Loránd University, Pázmány Péter sétány, 1/C, 1117 Budapest, Hungary. E-mail: gyuszk@cs.elte.hu. Supported by the Hungarian National Foundation for Scientific Research (OTKA) grant K60802.

S -paths having a high value and a low cost. Motivated by this, we consider the μ -weighted minimum-cost edge-disjoint S -paths problem (μ -CEDP):

$$\begin{aligned} \mu\text{-CEDP:} \quad & \text{Maximize } \sum_{P \in \mathcal{P}} \mu(s_P, t_P) - a(P) \\ & \text{over all edge-disjoint sets } \mathcal{P} \text{ of } S\text{-paths in } G. \end{aligned}$$

Here s_P, t_P are the terminals which P connects. Let $\text{val}(\mu; G, a)$ denote the optimal value of μ -CEDP.

In general, μ -CEDP is NP-hard. To see this, let $S = \{s, t, s', t'\}$ and define weight $\mu_{2\text{com}}$ by $\mu_{2\text{com}}(u, v) := 1$ if $\{u, v\} = \{s, t\}$ or $\{s', t'\}$, and $\mu_{2\text{com}}(u, v) := 0$ otherwise. Then $\mu_{2\text{com}}$ -CEDP with $a = 0$ is the integer version of 2-commodity flow maximization problem [19]. This problem is shown to be NP-hard [10].

However, there is an instance of terminal weights μ such that μ -CEDP is tractable, admits a nice combinatorial min-max theorem, and is not reducible to minimum-cost (single commodity) flow problem. Mader [33] established the following combinatorial min-max relation for the case $\mu = 1$ and $a = 0$ (1 denotes the all one function and 0 denotes the all zero function):

$$\text{val}(1; G, 0) = \frac{1}{2} \min \left(\sum_{s \in S} |\delta X_s| - \kappa \right), \quad (1.1)$$

where the minimum is taken over all families of disjoint node subsets X_s ($s \in S$) with $s \in X_s$. Here δX denotes the set of edges joining X and $V \setminus X$, and κ denotes the number of connected components K in $G - \bigcup_{s \in S} X_s$ such that $|\delta V(K)|$ is odd. Mader [34] extended (1.1) to openly node-disjoint S -paths packing; see [41] for a short proof. Lovász [31] derived this theorem from matroid matching theory. Schrijver [42] described an explicit reduction to a *linear* matroid matching problem. Thus a maximum number of edge-disjoint S -paths can be found in polynomial time by a linear matroid matching algorithm by Lovász [30]. In the 90's, Karzanov [23, 25] studied μ -CEDP for $\mu = p1$ ($p > 0$) and an arbitrary cost a ; if parameter p is large, then any optimal solution of this problem is a maximum edge-disjoint set of S -paths with the minimum total cost. He gave a min-max theorem and a combinatorial polynomial time algorithm for this problem. A proof outline was given in [25]. However the full proof [23] takes over 60 pages, is rather complicated, and is not submitted to any journal.

Our main result is a complete characterization of terminal weights μ for which μ -CEDP is tractable and admits a combinatorial min-max theorem, extending the edge-disjoint S -paths theorem by Mader and its min-cost generalization by Karzanov. To state the main result, let us introduce some notions. A *tree metric* is a weight function that can be represented as distances among points in a weighted tree. More precisely, a weight μ is called a tree metric if there exist a tree Γ , a nonnegative edge-length l , and a family of nodes $\{p_s\}_{s \in S} \subseteq V(\Gamma)$ indexed by S such that $\mu(s, t) = d_{\Gamma, l}(p_s, p_t)$ for $s, t \in S$, where $d_{\Gamma, l}$ denotes the shortest path metric of Γ with respect to edge-length l . By edge-subdivision, we can assume that l is uniform, i.e., $l(e) = \gamma (> 0)$ for all edges e . In this case, $d_{\Gamma, l}$ is also denoted by $d_{\Gamma, \gamma}$. Triple $(\Gamma, \{p_s\}_{s \in S}; \gamma)$ is called a *tree realization* of μ . Here γ is always taken as $1/2$ if μ is an integer-valued tree metric. Indeed, suppose (first) that μ

is represented by $\Gamma, \{p_s\}_{s \in S}, l$, as above. We can assume that for each vertex v of degree at most 2 there is $s \in S$ with $v = p_s$. Then each edge-length $l(e)$ is half-integral since $l(e)$ is equal to $(\mu(s, u) + \mu(t, v) + \mu(s, v) + \mu(t, u) - \mu(s, t) - \mu(u, v))/2$ for some $s, t, u, v \in S$. Hence, by edge-subdivision, we get a realization of μ with $\gamma = 1/2$.

Let $G = (V, E)$ be a graph with terminal set S . A node in $V \setminus S$ is called an *inner node*. G is said to be *inner Eulerian* if every inner node has an even degree. An *inner-odd-join* is an edge subset $F \subseteq E$ for which $G - F = (V, E \setminus F)$ is inner Eulerian. For a tree Γ with edge-length l and a map $\rho : V \rightarrow V(\Gamma)$, let $d_{\Gamma, l}^\rho$ denote a function on E defined by $d_{\Gamma, l}^\rho(e) = d_{\Gamma, l}(\rho(x), \rho(y))$ for $e = xy$. Now we are ready to state our main theorem:

Theorem 1.1. *Let μ be an integer-valued tree metric realized by $(\Gamma, \{p_s\}_{s \in S}; 1/2)$. Then we have*

$$\text{val}(\mu; G, a) = \min_{\rho} \max_F (d_{\Gamma, 1/2}^\rho - a)(E \setminus F), \quad (1.2)$$

where the maximum is taken over inner-odd-joins F and the minimum is taken over maps $\rho : V \rightarrow V(\Gamma)$ satisfying $\rho(s) = p_s$ for $s \in S$. Moreover there exists a polynomial time algorithm to find edge-disjoint S -paths attaining the maximum.

Relation to Mader's min-max formula. Mader's formula (1.1) can be deduced from our formula (1.2) as follows. First, observe that, in (1.1), the minimum is always attained by $\{X_s\}$ with the following property:

$$(1.3) \quad \text{Each } x \in X_s \text{ is reachable from } s \text{ in the subgraph of } G \text{ induced by } X_s.$$

Indeed, delete such nonreachable nodes from X_s . Then $\sum |\delta X_s| - \kappa$ does not increase. Next consider our formula (1.2). Let $\mu = 1$, and let Γ be the star with $|S|$ leaves p_s ($s \in S$) and center p_0 . Then $(\Gamma, \{p_s\}_{s \in S}; 1/2)$ realizes μ . A map $\rho : V \rightarrow V(\Gamma)$ satisfying $\rho(s) = p_s$ is identified with a set $\{X_s\}_{s \in S}$ of disjoint sets with $s \in X_s$ by correspondence $X_s \leftrightarrow \rho^{-1}(p_s)$. Then $d_{\Gamma, 1/2}^\rho(E) = \sum_{s \in S} |\delta X_s|/2$. Also in (1.2) the minimum is always attained by a map ρ with the property (1.3) (under the correspondence $X_s \leftrightarrow \rho^{-1}(p_s)$). To see this, consider a map ρ' obtained from ρ by replacing $\rho(x)$ with p_0 for all such nonreachable nodes x . Then the inner maximum does not increase. Indeed, take an inner-odd-join F' attaining $\max_F d_{\Gamma, 1/2}^{\rho'}(E \setminus F)$. Since $d_{\Gamma, 1/2}^{\rho'}(e) \leq d_{\Gamma, 1/2}^\rho(e)$ ($e \in E$) by the construction, we have $d_{\Gamma, 1/2}^{\rho'}(E \setminus F') \leq d_{\Gamma, 1/2}^\rho(E \setminus F') \leq \max_F d_{\Gamma, 1/2}^\rho(E \setminus F)$. Next we estimate $\min_F d_{\Gamma, 1/2}^\rho(F)$ under property (1.3). Let Y_1, Y_2, \dots, Y_m be the node sets of connected components of $G - \bigcup_{s \in S} X_s$. Any inner-odd-join F meets at least one edge in δY_i if $|\delta Y_i|$ is odd; hence $\min_F d_{\Gamma, 1/2}^\rho(F) \geq \kappa/2$. Conversely, taking one edge from each δY_i with $|\delta Y_i|$ odd, we obtain an edge set F with $d_{\Gamma, 1/2}^\rho(F) = \kappa/2$. By (1.3), we can greedily add edges e with $d_{\Gamma, 1/2}^\rho(e) = 0$ to F , and extend F to an inner-odd-join of G . This means $\min_F d_{\Gamma, 1/2}^\rho(F) = \kappa/2$. Hence Theorem 1.1 implies Mader's min-max formula (1.1).

Dual half-integrality of min-cost edge-disjoint S -paths problem. As mentioned already, our result includes Karzanov's min-cost edge-disjoint S -paths theo-

rem for a special case of $\mu = p1$. Therefore this paper includes a relatively simpler proof of his result. Furthermore our min-max formula includes half-integrality of dual. Indeed, it will turn out that the inner maximization is a minimum-cost T -join problem. Although edge-cost $c := d_{\Gamma,1/2^p} - a$ is half-integral, the sum of $c(e)$ along every cycle is an integer (since tree Γ is bipartite). By Seymour's odd-cut packing theorem [44] for bipartite graphs, the inner maximization is dualized to a half-integral odd-cut packing problem. This dual half-integrality affirmatively resolves a question raised by Karzanov [25, Section 6, (2)] as the special case; see Section 5 for detail.

Tractability classification of terminals weights. By considering $a = 0$ we get an important special case μ -EDP of μ -CEDP:

$$\mu\text{-EDP:} \quad \text{Maximize } \sum_{P \in \mathcal{P}} \mu(s_P, t_P) \text{ over all edge-disjoint sets } \mathcal{P} \text{ of } S\text{-paths in } G.$$

A terminal weight μ is said to be *EDP-tractable* if μ -EDP is solvable in polynomial time for every graph containing terminal set S . The all-one weight 1 is EDP-tractable. By our result, every tree metric is EDP-tractable. However, a slightly more general class of weights is EDP-tractable. A weight μ is said to be a *truncated tree metric* if it can be represented as distances between balls in a weighted tree, or equivalently, if there are a tree metric $\bar{\mu}$ and a nonnegative function (*radius*) r on S such that

$$\mu(s, t) = \max(\bar{\mu}(s, t) - r(s) - r(t), 0) \quad (s, t \in S). \quad (1.4)$$

Corollary 1.2. *Every truncated tree metric is EDP-tractable.*

Indeed, suppose that μ is a truncated tree metric represented as (1.4) for a tree metric $\bar{\mu}$ and radius r . Then μ -EDP reduces to $\bar{\mu}$ -CEDP as follows. First, for each terminal s , add a new inner node s' , and replace each edge sx incident to s by two edges ss' and $s'x$. Clearly the problem does not change. Next define cost a as: $a(e) = r(s)$ if $e = ss'$ for a terminal s and $a(e) = 0$ otherwise. Consider $\bar{\mu}$ -CEDP for this graph with this cost. Then value $\bar{\mu}(s, t) - a(P)$ of any (s, t) -path P is equal to $\bar{\mu}(s, t) - a(s) - a(t) (\leq \mu(s, t))$. Therefore the optimal value of the $\bar{\mu}$ -CEDP is at most that of the original μ -EDP. Conversely, from an optimal solution in the μ -EDP, we can delete all paths P with $\mu(s_P, t_P) = 0$ to obtain a solution in the $\bar{\mu}$ -CEDP of the same objective value. Note that the same idea also reduces μ -CEDP to $\bar{\mu}$ -CEDP.

We further prove that this result is best possible in the following sense:

Theorem 1.3. *If μ is not a truncated tree metric, then μ -EDP is NP-hard.*

This hardness result implies that EDP-tractable weights are exactly truncated tree metrics unless $P = NP$.

Application: Approximating edge-disjoint S -paths for “near” tree metric. Finding a good approximation for μ -EDP (with general μ) is a great challenge in the area of approximation algorithms [45]; the current best is an $O(\sqrt{|V|})$ -approximation due to Chekuri, Khanna and Shepherd [4]. The following observation suggests a possible use of our result to the design of approximation algorithms for μ -EDP:

For a weight μ , if we can choose some EDP-tractable weight μ^* such that

$$\mu^* \leq \mu \leq \alpha\mu^*$$

for some $\alpha \geq 1$, then any optimal solution for μ^* -EDP is an α -approximate solution for μ -EDP.

Here an α -approximate solution is a solution having the objective value at least $1/\alpha$ times the optimal value. In the case of metric weights, this problem—finding a good embedding into a tree metric—falls into the area of the *low-distortion embedding*; see [32]. For two metric spaces (X, d) and (Y, d') , a map $\phi : X \rightarrow Y$ is an *embedding with distortion α* if $d'(\phi(x), \phi(y)) \leq d(x, y) \leq \alpha d'(\phi(x), \phi(y))$ for $x, y \in X$. Given a metric μ (metric space (S, μ)), let α^* denote the minimum distortion of an embedding into a tree metric (the metric space on a tree). Finding a tree metric with minimum distortion α^* is known to be NP-hard. In the case where the input metric μ is a graph metric, Bădoiu, Indyk, and Sidiropoulos [2] gave a polynomial time algorithm to find an embedding into some tree metric with distortion $O(\alpha^*)$. Consequently, we have the following.

Corollary 1.4. *Let μ be a graph metric with the minimum distortion α^* into tree metrics. Then there exists an $O(\alpha^*)$ -approximation algorithm for μ -EDP.*

Later, Chepoi, Dragan, Newman, Rabinovich, and Vaxes [6] gave a polynomial time algorithm to find a tree embedding with distortion $6\alpha^*$. So there exists a $6\alpha^*$ -approximation algorithm for μ -EDP when μ is a graph metric with minimum distortion α^* into tree metrics. Unfortunately, this approach does not improve the approximation ratio (for general graph metrics) since the minimum distortion of general metrics into tree metrics is $\Omega(|S|)$ [39]. Also Bartal’s probabilistic embedding [1] does not work (since our problem is a “maximization” problem). Nevertheless we hope that our results and techniques will provide tools to approximation algorithms for μ -EDP/CEDP.

Multicommodity flows. A natural fractional relaxation of μ -EDP— μ -weighted maximum multiflow problem—has been well-studied in the literature. Hu [19] proved the existence of a *half-integral* optimal solution for the 2-commodity flow maximization problem (the case of $\mu = \mu_{2\text{com}}$). Lovász [29] and Cherkassky [5] independently proved the existence of a half-integral optimal solution for all-one weight $\mu = 1$. However the 3-commodity flow does not have such a property. Such phenomena of the (non-)existence of an optimal solution with bounded denominator lead to another kind of a weight classification problem, called the *fractionality problem*. The fractionality problem [21, 22] asks the classification of weights μ for which every μ -weighted multiflow problem has a $1/k$ -integral optimal multiflow for some integer k . Recent works [15, 16, 17] by Hirai gave a solution to the fractionality problem. He [18] also gave a solution to the *node-capacitated* variation as: μ has bounded fractionality for node-capacitated μ -weighted maximum multiflow problems if and only if μ can be represented as the distance between *subtrees* (not necessarily balls!) in a tree. This work is influenced by these developments.

Discrete minimax relation. The formula (1.2) can be seen as a discrete minimax relation. It is known in [26] that if G is inner Eulerian and μ is an integer-valued tree metric realized by $(\Gamma, \{p_s\}_{s \in S}; 1/2)$, then the μ -weighted maximum (fractional) multiflow problem has an integral optimal multiflow with value

$$\min_{\rho} d_{\Gamma, 1/2}^{\rho}(E),$$

where the minimum is taken over all maps $\rho : V \rightarrow V(\Gamma)$ with $\rho(s) = p_s$ for $s \in S$; see Section 2.1. Since the complement of the union of edge-disjoint S -paths is an inner-odd-join, for a graph $G = (V, E)$ (not necessarily inner Eulerian) we have

$$\text{val}(\mu; G, a) = \max_F (\text{val}(\mu; G - F, 0) - a(E \setminus F)) = \max_F \min_{\rho} (d_{\Gamma, 1/2}^{\rho} - a)(E \setminus F).$$

Hence our formula (1.2) implies a discrete minimax theorem stating that “*max*” and “*min*” are interchangeable. In particular, the weak duality of (1.2) can be seen by the general relation $\max_x \min_y f(x, y) \leq \min_y \max_x f(x, y)$.

Organization. The first goal is the proof of Theorem 1.1, and the second goal is the proof of Theorem 1.3. We construct an LP-relaxation of μ -CEDP/EDP according to the observation: the union of S -paths is an inner Eulerian graph, and hence the complement is an inner-odd-join. Thus $\text{val}(\mu; G, a) = \max_F \text{val}(\mu; G - F, 0) - a(E \setminus F)$ holds. Next relax inner-odd-joins into points in the convex hull of the incidence vectors of inner-odd-joins (*inner-odd-join polytope*), and relax edge-disjoint paths into *multiflows* in a natural way. This LP-relaxation was used in [3, 25, 27] (implicitly or explicitly). Note that the inner-odd-join polytope is a T -join polytope, and therefore a polynomial time membership oracle exists; see, e.g., [35]. Hence this LP-relaxation can be solved in polynomial time by the ellipsoid method [12]. We prove that if μ is a tree metric, then this LP-relaxation has an integral optimal solution. This extends a result of Keijsper, Pendavingh, and Stougie [27] for $\mu = 1$ and $a = 0$. Although their proof assumes Mader’s theorem (1.1), our proof does not, and is *purely polyhedral*, i.e., it consists only of LP-based techniques. So our paper includes a new proof of Mader’s edge-disjoint path theorem; it seems the first polyhedral proof.

In Section 2, we give preliminary arguments including the polyhedral description of inner-odd-join polytopes. In Section 3, we prove the first part of Theorem 1.1, which consists of three steps. The first step is to represent the dual of the LP-relaxation as a *facility location problem on a tree*. The second step is to show that a dual optimum is always attained by a nice structured combinatorial solution, and the optimal value coincides with the right hand side in the min-max formula (1.2). In the third step, we analyze the stability of dual combinatorial solutions under edge deletion; we see an unexpected application of the *polynomial uncrossing process* [25]. We then prove by induction the existence of an integral optimal solution. In Section 4, we prove the polynomial solvability (the second part of Theorem 1.1) and NP-hardness (Theorem 1.3). The polynomial solvability is almost proved by the argument in Section 3. The proof of the NP-hardness result is based on an interesting connection between a reduction to an integer 2-commodity flow feasibility problem and 4-point characterization of truncated tree metrics. As is well known,

a metric μ is a tree metric if and only if there is no quadruple (s, t, s', t') (*violating quadruple*) satisfying $\mu(s, t) + \mu(s', t') > \max\{\mu(s, s') + \mu(t, t'), \mu(s', t) + \mu(s, t')\}$; see [43]. We prove a similar 4-point characterization for truncated tree metrics. From a violating quadruple (s, t, s', t') , we can reduce a version of the integer 2-commodity flow feasibility problem, which is NP-hard, to $\mu|_{S'}$ -EDP on the four terminal $S' = \{s, t, s', t'\}$. In Section 5, we discuss related issues and raise future research directions.

Notation. Let \mathbf{R} and \mathbf{R}_+ denote the sets of reals and nonnegative reals, respectively, and let \mathbf{Z} and \mathbf{Z}_+ denote the sets of integers and nonnegative integers, respectively. For a set X , let \mathbf{R}^X and \mathbf{R}_+^X denote the sets of functions from X to \mathbf{R} and from X to \mathbf{R}_+ , respectively. For a function $f : X \rightarrow \mathbf{R}$, let $\mathcal{D}^+ f$ and $\mathcal{D}^- f$ denote the sets of elements $x \in X$ with $f(x) > 0$ and $f(x) < 0$, respectively. Let f^+ , f^- , and $|f|$ denote the functions defined by $f^+(x) = \max(0, f(x))$, $f^-(x) = \min(0, f(x))$, and $|f|(x) = \max(-f(x), f(x))$, respectively. Let $\|f\|$ denote the l_∞ -norm of f ; $\|f\| := \max_{x \in X} |f(x)|$. For a subset $Y \subseteq X$, let $f(Y)$ denote the sum of $f(y)$ over all $y \in Y$. The all-one function is denoted by 1, and the zero function is denoted by 0.

By a *metric* on a set V , we mean a function $d : V \times V \rightarrow \mathbf{R}_+$ satisfying $d(x, x) = 0$, $d(x, y) = d(y, x)$, and triangle inequalities $d(x, y) + d(y, z) \geq d(x, z)$ for $x, y, z \in V$. For a subset $X \subseteq V$, the *cut metric* δ_X is defined by $\delta_X(x, y) = 1$ if $|X \cap \{x, y\}| = 1$ and $\delta_X(x, y) = 0$ otherwise.

Let $G = (V, E)$ be a graph with possible parallel edges and loops. For an edge e , we write $e = xy$ if e joins nodes x, y . For a node subset $X \subseteq V$, let $\delta X = \delta^G X$ denote the set of edges joining X and $V \setminus X$. For two functions f, g on E , let $f \cdot g$ denote the inner product $\sum_{e \in E} f(e)g(e)$. We regard a metric d on V as a function on E by $d(e) := d(x, y)$ for $e = xy \in E$. In particular, $\delta_X \cdot \xi = \xi(\delta X)$ for $\xi \in \mathbf{R}^E$.

2 Preliminaries

Let $G = (V, E)$ be an undirected graph with terminal set $S \subseteq V$, μ a terminal weight, and a an edge-cost. Recall that an inner-odd-join is an edge subset F for which $G - F$ is inner Eulerian. Our starting point is the following relation:

$$\text{val}(\mu; G, a) = \max_F (\text{val}(\mu; G - F, 0) - a(E \setminus F)), \quad (2.1)$$

where the maximum is taken over all inner-odd-joins F in G . To see this, let \mathcal{P}^* be an edge-disjoint set of S -paths attaining $\text{val}(\mu; G, a)$, and let F^* be the complement of the union of \mathcal{P}^* . Then F^* is an inner-odd-join, and $\text{val}(\mu; G, a) = \sum_{P \in \mathcal{P}^*} \mu(s_P, t_P) - a(E \setminus F^*) \leq \text{val}(\mu; G - F^*, 0) - a(E \setminus F^*)$. Conversely, for any inner-odd-join F and any edge-disjoint set \mathcal{P} of S -paths in $G - F$, we have $\sum_{P \in \mathcal{P}} \mu(s_P, t_P) - a(E \setminus F) \leq \sum_{P \in \mathcal{P}} \mu(s_P, t_P) - a(P) \leq \text{val}(\mu; G, a)$, where we use the nonnegativity of a in the first inequality.

Motivated by (2.1), we consider the following linear programming relaxation:

$$\begin{aligned} \text{Max.} \quad & \sum_{P \in \Pi} \mu(s_P, t_P) f(P) - a(E) + \sum_{e \in E} a(e) \xi(e) & (2.2) \\ \text{s. t.} \quad & \xi(e) + \sum_{P \in \Pi: e \in P} f(P) \leq 1 \quad (e \in E), \\ & \xi \in Q, \quad f : \Pi \rightarrow \mathbf{R}_+. \end{aligned}$$

Here $Q = Q_{G,S} \subseteq \mathbf{R}_+^E$ denotes the convex hull of the incidence vectors of all inner-odd-joins in G , and $\Pi = \Pi_{G,S}$ denotes the set of all S -paths in G . The maximum value of (2.2) is denoted by $\text{val}(\mu; G, a)$. Obviously $\text{val}(\mu; G, a) \leq \overline{\text{val}}(\mu; G, a)$. We are going to show that if μ is a tree metric, then (2.2) has an integral optimal solution.

2.1 Laminar locking theorem

In the case where G is inner Eulerian and $a = 0$, our problem becomes easier; we may put $\xi = 0$ in (2.2). So (2.2) coincides with the natural LP-relaxation of μ -EDP. We will use the following fact as the base case of our inductive argument:

Lemma 2.1. *If μ is a tree metric, G is inner Eulerian, and $a = 0$, then there exists an integral optimal solution (ξ^*, f^*) in (2.2) with $\xi^* = 0$.*

This is a special case of the *multiflow locking theorem* due to Karzanov-Lomonosov [26]; see [42, Section 77.3c]. The multiflow locking theorem says that if a family $\mathcal{F} \subseteq 2^S$ has no pairwise crossing triple and G is inner Eulerian, then there exists an integral multiflow f being simultaneously a maximum (single-commodity) $(A, S \setminus A)$ -flow for all $A \in \mathcal{F}$. Here a pair of sets $X, Y \subseteq S$ is said to be *crossing* if $X \cap Y$, $X \setminus Y$, $Y \setminus X$, and $S \setminus (X \cup Y)$ are all nonempty. One can easily see that such a multiflow f is a maximum multiflow with respect to a terminal weight $\mu = \sum_{X \in \mathcal{F}} \beta(X) \delta_X$ for any $\beta : \mathcal{F} \rightarrow \mathbf{R}_+$. As is well-known, a tree metric is just a nonnegative sum of cut metrics for a family without crossing pairs; see Section 2.2 below.

A simpler and shorter proof of the multiflow locking theorem is available at [11], and a faster algorithm finding an integer optimal solution is given by [20].

2.2 Trees and laminar families

A *metric-tree* \mathcal{T} is a metric space isometric to a 1-dimensional contractible complex endowed with the length metric. The metric function is denoted by $d_{\mathcal{T}}$. For a point $p \in \mathcal{T}$ and nonnegative $\epsilon \geq 0$, the closed ball $B(p, \epsilon)$ and the open ball $B^\circ(p, \epsilon)$ are defined as the subsets of points $q \in \mathcal{T}$ with $d_{\mathcal{T}}(p, q) \leq \epsilon$ and with $d_{\mathcal{T}}(p, q) < \epsilon$, respectively. For two points p, q , the closed interval $[p, q]$ is defined as the set of points t with $d_{\mathcal{T}}(p, q) = d_{\mathcal{T}}(p, t) + d_{\mathcal{T}}(t, q)$. The open interval (p, q) is defined as $[p, q] \setminus \{p, q\}$. A *vertex* of \mathcal{T} is a point q such that, for every $\epsilon > 0$, the open ball $B^\circ(q, \epsilon)$ is not an open interval. In the case where \mathcal{T} is endowed with a special point $r \in \mathcal{T}$, called a *root*, we can define a partial order \preceq by $p \preceq q \Leftrightarrow [r, p] \subseteq [r, q]$. By $p \prec q$ we mean $[r, p] \subset [r, q]$ (proper inclusion). A metric-tree of two vertices is called a *segment*.

Let Γ be a tree (in the graph-theoretical sense). From Γ we can construct a metric-tree as follows. For each edge $e = uv \in E(\Gamma)$, consider segment F_e with two vertices $p_{e,u}, p_{e,v}$ and unit length $d_{F_e}(p_{e,u}, p_{e,v}) = 1$. Then consider the disjoint union $\bigcup_{e \in E} F_e$, and for each node $u \in V(\Gamma)$, identify points $p_{e,u}$ for all edges e incident to u ; the image of $p_{e,u}$ is denoted by p_u . The resulting metric-tree is denoted by $\bar{\Gamma}$. Then $(V(\Gamma), d_\Gamma)$ isometrically embeds into $(\bar{\Gamma}, d_{\bar{\Gamma}})$ by $v \mapsto p_v$. Hence we can identify v and p_v , and can assume $V(\Gamma) \subseteq \bar{\Gamma}$.

Recall the tree representation of a laminar family; see [28, Section 2.2] and [42, Section 13.4]. A pair of two sets $X, Y \subseteq V$ is said to be *intersecting* if $X \cap Y$, $X \setminus Y$, and $Y \setminus X$ are all nonempty. A family of sets without intersecting pairs is called *laminar*.

Let φ be a map from V to a metric-tree \mathcal{T} . Let $d_{\mathcal{T}^\varphi}$ denote the metric on V defined by $d_{\mathcal{T}^\varphi}(x, y) := d_{\mathcal{T}}(\varphi(x), \varphi(y))$ for $x, y \in V$. Let $\mathcal{E}_{\mathcal{T}, \varphi}$ be the set of inclusion-maximal connected subsets $\omega \subseteq \mathcal{T}$ not meeting the image of φ , vertices, and root (if specified). Each $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$ is an open segment. Let π_ω denote its length. Suppose that \mathcal{T} has a root r . Let X_ω denote the set of nodes $x \in V$ such that $\varphi(x)$ belongs to the connected component of $\mathcal{T} \setminus \{\omega\}$ not containing r . Then one can easily see that

$$(2.3) \quad \{X_\omega \mid \omega \in \mathcal{E}_{\mathcal{T}, \varphi}\} \text{ is laminar, and } d_{\mathcal{T}^\varphi} = \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi}} \pi_\omega \delta_{X_\omega}.$$

Conversely, for any set function $\pi : 2^V \rightarrow \mathbf{R}_+$ having a laminar support, there exist a metric-tree \mathcal{T} with root r and a map $\varphi : V \rightarrow \mathcal{T}$ such that $\{X_\omega \mid \omega \in \mathcal{E}_{\mathcal{T}, \varphi}\} = \mathcal{D}^+ \pi$ and $\pi_\omega = \pi(X_\omega)$ for $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$; see [43]. In this case, we say that (\mathcal{T}, φ) *realizes* π . In particular, a tree metric is a nonnegative sum of cut metrics for a laminar family.

2.3 T -joins and inner-odd-joins

Let $G = (V, E)$ be an undirected graph, and let T be a node subset $T \subseteq V$ having an even cardinality. For an edge subset $F \subseteq E$, let $\text{odd}(F)$ denote the set of nodes incident to an odd number of edges in F . An edge subset F is called a *T -join* if $\text{odd}(F) = T$. See [28, Chapter 12] and [42, Chapter 29] for basics on T -joins. Let S be a terminal subset of V . An inner-odd-join is an edge subset F such that $G - F$ is inner-Eulerian. Let G/S be the graph obtained from G by identifying S into one node. Hence $E(G/S)$ is naturally identified with E . We observe the following relation between T -joins and inner-odd-joins.

Lemma 2.2. *F is an inner-odd-join in G with terminal set S if and only if F is an $\text{odd}(E)$ -join in G/S .*

Thus any argument for inner-odd-joins falls into theory of T -joins. We are interested in the minimum cost inner-odd-join problem with a cost vector $w : E \rightarrow \mathbf{R}$. Let $h[w] = h^G[w]$ be a set function on $V \setminus S$ defined by

$$h[w](X) := \begin{cases} 1 & \text{if } |\delta X \setminus \mathcal{D}^- w| \text{ is odd,} \\ 0 & \text{if } |\delta X \setminus \mathcal{D}^- w| \text{ is even,} \end{cases} \quad (X \subseteq V \setminus S).$$

Lemma 2.3. *For $w : E \rightarrow \mathbf{R}$, the following values are equal:*

- (1) $\min\{w(F) \mid F: \text{inner-odd-join}\}$.
- (2) $w^-(E) + \min\{|w| \cdot \xi \mid \xi \in \mathbf{R}_+^E, \delta_X \cdot \xi \geq h[w](X) (X \subseteq V \setminus S)\}$.
- (3) $w^-(E) + \max\left\{\sum_X \pi(X)h[w](X) \mid \sum_X \pi(X)\delta_X \leq |w|, \pi : 2^{V \setminus S} \rightarrow \mathbf{R}_+\right\}$.

Proof. (2) \equiv (3) follows from LP-duality. We show (1) \equiv (2). By a well-known trick ([42, Section 29.1] and [28, Section 12.2]), we first reduce (1) to a minimum cost T -join problem for a *nonnegative* cost vector $|w|$. Observe that, for $F, H \subseteq E$, symmetric difference $H \Delta F$ is an odd(E)-join if and only if F is an odd($E \setminus H$)-join. Therefore, by taking \mathcal{D}^-w as H , we obtain

$$\begin{aligned}
\min\{w(F) \mid F: \text{inner-odd-join}\} &= \min\{w(F) \mid F: \text{odd}(E)\text{-join in } G/S\} \quad (2.4) \\
&= \min\{w(F' \Delta \mathcal{D}^-w) \mid F': \text{odd}(E \setminus \mathcal{D}^-w)\text{-join in } G/S\} \\
&= \min\{w(F') + w(\mathcal{D}^-w) - 2w(F' \cap \mathcal{D}^-w) \mid F': \text{odd}(E \setminus \mathcal{D}^-w)\text{-join in } G/S\} \\
&= w^-(E) + \min\{|w|(F') \mid F': \text{odd}(E \setminus \mathcal{D}^-w)\text{-join in } G/S\}.
\end{aligned}$$

Since $|w|$ is nonnegative, the second term of the last equation is equivalent to a linear minimization over the uphull $P + \mathbf{R}_+^E$ of the T -join polytope P for G/S with $T := \text{odd}(E \setminus \mathcal{D}^-w)$. By Edmonds-Johnson theorem [9], the uphull $P + \mathbf{R}^E$ is the set of points ξ satisfying

$$\begin{aligned}
\xi(e) &\geq 0 & (e \in E), \\
\xi(\delta^{G/S}X) &\geq 1 & (X \subseteq V(G/S), |X \cap T| : \text{odd}).
\end{aligned} \quad (2.5)$$

A node subset in G/S is identified with a node subset X in G with $X \subseteq V \setminus S$ or $S \subseteq X$; clearly $\delta^G X = \delta^{G/S} X$. Moreover, for $H \subseteq E$, $|X \cap \text{odd}(H)|$ is odd if and only if $|\delta X \cap H|$ is odd. By substituting (2.5) to the last equation in (2.4), and using $h[w]$, we obtain (1) \equiv (2). \square

The explicit inequality description of inner-odd-join polytope Q is also given as

$$\begin{aligned}
0 &\leq \xi(e) \leq 1 & (e \in E), \\
\xi(\delta X \setminus F) + (1 - \xi)(\delta X \cap F) &\geq h[-1_F](X) & (X \subseteq V \setminus S, F \subseteq E),
\end{aligned} \quad (2.6)$$

where 1_F denotes the incidence vector of F . This is also a consequence of Lemma 2.2 and Edmond-Johnson theorem.

We end this subsection with noting the following fundamental properties of $h = h[w]$, which immediately follow from the definition.

Lemma 2.4. *For $X, Y \subseteq V \setminus S$, we have the following.*

- (1) *If $X \cap Y = \emptyset$, then $h(X \cup Y) \equiv h(X) + h(Y) \pmod{2}$.*
- (2) *If $h(X) = h(Y) = 1$, then $h(X \cap Y) = h(X \cup Y) = 1$ or $h(X \setminus Y) = h(Y \setminus X) = 1$.*

2.4 Polynomial uncrossing process

In this subsection, we study the following cut packing program arising from (3) in Lemma 2.3, where we denote $h[w]$ by h .

$$\text{Max. } \sum_X \pi(X) \quad \text{s.t. } \sum_X \pi(X)\delta_X \leq |w|, \quad \pi : \mathcal{D}^+h \rightarrow \mathbf{R}_+. \quad (2.7)$$

We are interested in a feasible solution π of (2.7) such that support $\mathcal{D}^+\pi$ is laminar. Such a feasible solution is called *laminar*.

Now we are given an arbitrary feasible function π of (2.7). Consider the following process, called an *uncrossing process*.

step 0: If $\mathcal{D}^+\pi$ is laminar, then stop.

step 1: Choose an intersecting pair $X, Y \in \mathcal{D}^+\pi$ and let $\alpha := \min(\pi(X), \pi(Y))$.

step 2: Choose $(X', Y') \in \{(X \cup Y, X \cap Y), (X \setminus Y, Y \setminus X)\}$ with $X', Y' \in \mathcal{D}^+h$, decrease π by α on X, Y and increase π by α on X', Y' . Go to **step 0**.

Here the existence of X', Y' in **step 2** is guaranteed by Lemma 2.4 (2). The operation in **step 2** is called an *uncrossing*. By an arbitrary choice of X, Y, X', Y' in each iteration, this process terminates after finitely many iterations. So we can make any feasible solution π laminar, keeping the feasibility and the objective value. In particular the maximum of (2.7) is always attained by a laminar solution.

In [24], Karzanov proved that, by an appropriate choice of X, Y, X', Y' , the uncrossing process for π terminates so that the number of uncrossings is bounded by a polynomial of $|V|$ and $|\mathcal{D}^+\pi| \leq |2^V|$. The important point is that this bound does not depend on the bit size representing the numerical values of π . Hence we have the following.

Theorem 2.5 ([24]). *There exists a function $p : \mathbf{Z}_+ \rightarrow \mathbf{Z}_+$ such that for every feasible solution π in (2.7) some uncrossing process to π terminates after at most $p(|V|)$ uncrossings.*

This result has an unexpected application about a *stability* of laminar feasible solutions in (2.7); recall that $\|\cdot\|$ is the l_∞ -norm.

Lemma 2.6. *Let π be a nonoptimal laminar feasible solution in (2.7). For every $\epsilon > 0$ there exists a laminar feasible solution π^* in (2.7) having a bigger objective value and satisfying $\|\pi^* - \pi\| < \epsilon$.*

Proof. The proof technique is important for us. Take sufficiently small $\epsilon' > 0$; more precisely, take ϵ' satisfying

$$0 < \epsilon' < \frac{\min(\epsilon, \min_{X \in \mathcal{D}^+\pi} \pi(X)/2)}{2^{p(|V|)+1} - 1}.$$

By convexity there exists a (not necessarily laminar) feasible solution π' having a bigger objective value and satisfying $\|\pi' - \pi\| < \epsilon'$. Next apply Karzanov's uncrossing process to π' . Then we get a laminar feasible solution π^* having a bigger objective value than that of π .

We show $\|\pi^* - \pi\| < \epsilon$. By $\epsilon' < \min_{X \in \mathcal{D}^+\pi} \pi(X)$, we have $\mathcal{D}^+\pi \subseteq \mathcal{D}^+\pi'$. Therefore each uncrossing step chooses X, Y so that at least one of X, Y does not belong to $\mathcal{D}^+\pi$. In each iteration α takes the value on $\mathcal{D}^+h \setminus \mathcal{D}^+\pi$, and the change in the uncrossing is bounded by the maximum value on $\mathcal{D}^+h \setminus \mathcal{D}^+\pi$. Moreover, by Theorem 2.5, the number of uncrossings is bounded by $p(|V|)$. From this we have

$$\|\pi - \pi^*\| \leq \|\pi - \pi'\| + \|\pi' - \pi^*\| \leq \epsilon' + \sum_{k=1}^{p(|V|)} 2^k \epsilon' = (2^{p(|V|)+1} - 1)\epsilon' < \epsilon. \quad (2.8)$$

□

3 Combinatorial min-max theorem

The goal of this section is to prove the min-max relation (1.2) in Theorem 1.1. Throughout this section, $G = (V, E)$ is an undirected graph with terminal set $S \subseteq V$, μ is a terminal weight on S , and a is an edge-cost. Since $\text{val}(2\mu; G, 2a) = 2 \text{val}(\mu; G, a)$, $\overline{\text{val}}(2\mu; G, 2a) = 2\overline{\text{val}}(\mu; G, a)$, and $d_{\Gamma, 1/2} = d_{\Gamma, 1}/2$, it suffices to consider the case where both μ and a are even-valued (by taking $(2\mu, 2a)$ as (μ, a)). Hence we may assume that μ is realized by $(\Gamma, \{p_s\}_{s \in S}; 1)$. Also we may assume the following nonredundancy:

$$(3.1) \quad \text{For each } p \in V(\Gamma) \text{ of degree one, there is } s \in S \text{ with } p = p_s.$$

In Section 3.1, we introduce a special dual of (2.2), which is a continuous relaxation of the RHS of (1.2). In Section 3.2, we show that the optimal value of this special dual is always attained by a discrete solution, which implies that the optimal value of (2.2) is equal to the RHS of (1.2). Finally, in Section 3.3, we prove that (2.2) has an integral optimal solution, which establishes the formula (1.2).

3.1 Duality and stability

We introduce a special dual problem of (2.2) and show that this problem has a kind of a stability property, similar to Lemma 2.6. We consider a triple $(\rho, \varphi, \mathcal{T})$ of a metric-tree \mathcal{T} with root r , maps $\rho : V \rightarrow \bar{\Gamma}$ and $\varphi : V \rightarrow \mathcal{T}$. Regard (ρ, φ) as a map from V to the Cartesian product $\bar{\Gamma} \times \mathcal{T}$ of two metric-trees.

A triple $(\rho, \varphi, \mathcal{T})$ is said to be *feasible* if it satisfies

$$(3.2) \quad (1) \quad (\rho, \varphi)(s) = (p_s, r) \text{ for } s \in S, \text{ and} \\ (2) \quad d_{\mathcal{T}^\varphi}(e) \leq |d_{\bar{\Gamma}^\rho} - a|(e) \text{ for } e \in E.$$

For a feasible triple $(\rho, \varphi, \mathcal{T})$, define $\tau(\rho, \varphi, \mathcal{T})$ by

$$\tau(\rho, \varphi, \mathcal{T}) := (d_{\bar{\Gamma}^\rho} - a)^+(E) - \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi}} \pi_\omega h[d_{\bar{\Gamma}^\rho} - a](X_\omega). \quad (3.3)$$

Define notion $h[d_{\bar{\Gamma}^\rho} - a] \circ (\varphi, \mathcal{T})$ by

$$h[d_{\bar{\Gamma}^\rho} - a] \circ (\varphi, \mathcal{T}) := \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi}} \pi_\omega h[d_{\bar{\Gamma}^\rho} - a](X_\omega). \quad (3.4)$$

Then τ is represented as

$$\tau(\rho, \varphi, \mathcal{T}) = (d_{\bar{\Gamma}^\rho} - a)^+(E) - h[d_{\bar{\Gamma}^\rho} - a] \circ (\varphi, \mathcal{T}).$$

A feasible triple $(\rho, \varphi, \mathcal{T})$ is said to be *rational* if the length of each segment in $\mathcal{E}_{\bar{\Gamma}, \rho}$ and in $\mathcal{E}_{\mathcal{T}, \varphi}$ is rational. The goal of this subsection is to prove the following two lemmas:

Lemma 3.1. *The following values are equal:*

- (1) $\overline{\text{val}}(\mu; G, a)$.
- (2) $\min \tau(\rho, \varphi, \mathcal{T})$, where the minimum is taken over all rational feasible triples $(\rho, \varphi, \mathcal{T})$.
- (3) $\min \max (d - a)(E \setminus F)$, where the maximum is taken over all inner-odd-joins F , and the minimum is taken over all metrics d on V satisfying $d(s, t) \geq \mu(s, t)$ for $s, t \in S$.
- (4) $\min \max (d_{\bar{\Gamma}^\rho} - a)(E \setminus F)$, where the maximum is taken over all inner-odd-joins F , and the minimum is taken over all maps $\rho : V \rightarrow \bar{\Gamma}$ with $\rho(s) = p_s$ for $s \in S$.
- (5) *The optimal value of the following problem:*

$$\begin{aligned} \text{Min.} \quad & (d - a)^+(E) - \sum_X \pi(X) h[d - a](X) & (3.5) \\ \text{s.t.} \quad & \sum_X \pi(X) \delta_X(e) \leq |d - a|(e) \quad (e \in E), \\ & d(s, t) \geq \mu(s, t) \quad (s, t \in S), \\ & d: \text{metric on } V, \\ & \pi : 2^{V \setminus S} \rightarrow \mathbf{R}_+. \end{aligned}$$

Note that (3.5) is a linear program if d is fixed.

Lemma 3.2 (Stability Lemma). *Let $(\rho, \varphi, \mathcal{T})$ be a nonoptimal feasible triple with $h[d_{\bar{\Gamma}^\rho} - a](X_\omega) = 1$ ($\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$). For every $\epsilon > 0$, there exists a rational feasible triple $(\rho', \varphi', \mathcal{T}')$ such that $\tau(\rho', \varphi', \mathcal{T}') < \tau(\rho, \varphi, \mathcal{T})$, $\|d_{\bar{\Gamma}^{\rho'}} - d_{\bar{\Gamma}^\rho}\| \leq \epsilon$, and $\|d_{\mathcal{T}' \varphi'} - d_{\mathcal{T} \varphi}\| \leq \epsilon$.*

Proof of Lemma 3.1. (1) \equiv (3). Dualize (2.2) with fixed $\xi \in Q$. Then we get

$$\overline{\text{val}}(\mu; G, a) = \max_{\xi \in Q} \min_l \{(l - a)(E) - (l - a) \cdot \xi\}, \quad (3.6)$$

where the minimum is taken over all $l : E \rightarrow \mathbf{R}_+$ satisfying $l(P) \geq \mu(s_P, t_P)$ for $P \in \Pi$. By the minimax theorem we obtain

$$\begin{aligned} \max_{\xi \in Q} \min_l \{(l - a)(E) - (l - a) \cdot \xi\} &= \min_l \max_{\xi \in Q} \{(l - a)(E) - (l - a) \cdot \xi\} \\ &= \min_l \max_F (l - a)(E \setminus F). \end{aligned}$$

Note that the first equality can also be directly obtained by the LP-duality with the help of the explicit description (2.6) of Q .

Let $g : \mathbf{R}_+^E \rightarrow \mathbf{R}$ be defined by

$$g(l) := \max_F (l - a)(E \setminus F) \quad (l \in \mathbf{R}_+^E), \quad (3.7)$$

where the maximum is taken over all inner-odd-joins F . Here g is monotone nondecreasing. Indeed, for $l \leq l'$, take an inner-odd-join F^* with $g(l) = (l - a)(E \setminus F^*)$. Then $(l - a)(E \setminus F^*) \leq (l' - a)(E \setminus F^*) \leq g(l')$. Therefore we can replace l by the shortest path metric $d_{G,l}$ on G with respect to l . From this we obtain (1) \equiv (3); obviously the minimum of g is attained by a rational solution.

(3) \equiv (4). (4) \geq (3) is obvious. Since g is monotone nondecreasing, (3) \geq (4) follows from the following claim; the proof is given for completeness.

Claim 3.3 ([16, 18]). For any metric d on V with $d(s, t) \geq \mu(s, t)$ for $s, t \in S$, there exists a map $\rho : V \rightarrow \bar{\Gamma}$ such that $\rho(s) = p_s$ ($s \in S$) and $d_{\bar{\Gamma}^\rho}(x, y) \leq d(x, y)$ for $x, y \in V$.

Proof. Suppose $V = \{x_1, x_2, \dots, x_n\}$ and $S = \{x_1, x_2, \dots, x_k\}$. Define $\rho : V \rightarrow \bar{\Gamma}$ recursively by

$$\rho(x_i) := \begin{cases} p_{x_i} & \text{if } i \leq k, \\ \text{an arbitrary point in } \bigcap_{j < i} B(\rho(x_j), d(x_j, x_i)) & \text{if } i > k, \end{cases} \quad (1 \leq i \leq n).$$

We show by induction that $\bigcap_{j < i} B(\rho(x_j), d(x_j, x_i))$ is nonempty, i.e., ρ is well-defined. If this is true, then $\rho(x_i) \in B(\rho(x_j), d(x_j, x_i))$ implies $d_{\bar{\Gamma}^\rho}(x_i, x_j) \leq d(x_i, x_j)$.

Recall that any collection of subtrees has the Helly property. So it suffices to show the *pairwise* nonempty intersection. Here two balls $B(p, r)$ and $B(p', r')$ intersect if and only if $d_{\bar{\Gamma}}(p, p') \leq r + r'$. Therefore $B(\rho(x_j), d(x_j, x_i)) \cap B(\rho(x_j), d(x_j, x_{j'})) \neq \emptyset$ follows from $d(x_i, x_j) + d(x_i, x_{j'}) \geq d(x_j, x_{j'}) \geq d_{\bar{\Gamma}^\rho}(x_j, x_{j'})$, where the last inequality follows from the induction. \square

In the proof, if d is rational, then we can choose ρ so that the length of each segment in $\mathcal{E}_{\bar{\Gamma}, \rho}$ is rational.

(3) \equiv (5). This follows from Lemma 2.3.

(5) \equiv (2). For a feasible solution (d, π) in (3.5), let $\tau(d, \pi)$ denote the objective value. For a feasible triple $(\rho, \varphi, \mathcal{T})$, let $\pi^{\varphi, \mathcal{T}} : 2^{V \setminus S} \rightarrow \mathbf{R}_+$ be defined by

$$\pi^{\varphi, \mathcal{T}}(X) = \begin{cases} \pi_\omega & \text{if } X = X_\omega \text{ for } \omega \in \mathcal{E}_{\mathcal{T}, \varphi}, \\ 0 & \text{otherwise,} \end{cases} \quad (X \subseteq V \setminus S). \quad (3.8)$$

By (2.3) and (3.2) (2), $(d_{\bar{\Gamma}^\rho}, \pi^{\varphi, \mathcal{T}})$ is feasible to (3.5) with $\tau(d_{\bar{\Gamma}^\rho}, \pi^{\varphi, \mathcal{T}}) = \tau(\rho, \varphi, \mathcal{T})$. Conversely, by (4) \equiv (5) we can take an optimal solution (d, π) in (3.5) such that $d = d_{\bar{\Gamma}^\rho}$ for a map $\rho : V \rightarrow \bar{\Gamma}$ with $\rho(s) = p_s$ for $s \in S$. By the uncrossing process in the previous section, we can make π laminar, with keeping $(d_{\bar{\Gamma}^\rho}, \pi)$ feasible (and optimal). There are a metric-tree \mathcal{T} with root r and a map $\varphi : V \rightarrow \mathcal{T}$ such that (φ, \mathcal{T}) realizes π . Then we have $d_{\mathcal{T}^\varphi}(e) = \sum_X \pi(X) \delta_X(e) \leq |d_{\bar{\Gamma}^\rho} - a|(e)$ for $e \in E$. Also $\varphi(S) = \{r\}$ by $\mathcal{D}^+ \pi \subseteq 2^{V \setminus S}$. This means that $(\rho, \varphi, \mathcal{T})$ is a feasible triple with $\tau(\rho, \varphi, \mathcal{T}) = \tau(d, \pi)$. We can assume that both d and π are rational-valued. Consequently $(\rho, \varphi, \mathcal{T})$ can be taken to be rational. The proof of Lemma 3.1 is complete. \square

Proof of Lemma 3.2. Suppose that $(\rho, \varphi, \mathcal{T})$ is a nonoptimal feasible triple with $h[d_{\bar{F}}^\rho - a](X_\omega) = 1$ for $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$. Our goal is to show the following claim:

Claim 3.4. For every $\tilde{\epsilon} > 0$, there exists a rational feasible solution $(\tilde{d}, \tilde{\pi})$ in (3.5) such that

- (1) $\tau(\tilde{d}, \tilde{\pi}) < \tau(\rho, \varphi, \mathcal{T})$,
- (2) $\|\tilde{d} - d_{\bar{F}}^\rho\| \leq \tilde{\epsilon}$,
- (3) $\|\tilde{\pi} - \pi^{\varphi, \mathcal{T}}\| \leq \tilde{\epsilon}$, and
- (4) $\tilde{d} = d_{\bar{F}}^{\tilde{\rho}}$ for a map $\tilde{\rho} : V \rightarrow V(\bar{F})$ with $\tilde{\rho}(s) = p_s$ ($s \in S$).

Recall that $\tau(d, \pi)$ is the objective value of (3.5), and $\pi^{\varphi, \mathcal{T}}$ is defined in (3.8). Suppose that this claim is true. Take a sufficiently small $\tilde{\epsilon} > 0$, and take $(\tilde{d}, \tilde{\pi})$ with properties (1-4) in this claim. By (2), $\mathcal{D}^\sigma(d_{\bar{F}}^\rho - a) \subseteq \mathcal{D}^\sigma(\tilde{d} - a)$ holds for $\sigma \in \{+, -\}$. By this fact together with $|d_{\bar{F}}^\rho - a|(e) \geq d_{\mathcal{T}^\varphi}(e) \geq \pi_\omega > 0$ ($\omega \in \mathcal{E}_{\mathcal{T}, \varphi}, e \in \delta X_\omega$), $d_{\bar{F}}^\rho - a$ and $\tilde{d} - a$ have the same sign pattern on δX_ω . This means $h[\tilde{d} - a](X_\omega) = h[d_{\bar{F}}^\rho - a](X_\omega) = 1$ ($\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$). Therefore we can assume

$$\mathcal{D}^+ \pi^{\varphi, \mathcal{T}} \subseteq \mathcal{D}^+ \tilde{\pi} \subseteq \mathcal{D}^+ h[\tilde{d} - a].$$

As in the proof of Lemma 2.6, applying Karzanov's uncrossing process to $\tilde{\pi}$ (with $w = \tilde{d} - a$), we obtain laminar $\tilde{\pi}'$ such that $(\tilde{d}, \tilde{\pi}')$ is (rational) feasible in (3.5), $\tau(\tilde{d}, \tilde{\pi}') \leq \tau(d', \pi')$, and $\|\tilde{\pi}' - \pi^{\varphi, \mathcal{T}}\| \leq C\epsilon'$, where C is a constant independent of ϵ' ; see (2.8). Take $(\tilde{\varphi}, \tilde{\mathcal{T}})$ realizing $\tilde{\pi}'$. Then $\tau(\tilde{\rho}, \tilde{\varphi}, \tilde{\mathcal{T}}) < \tau(\rho, \varphi, \mathcal{T})$, $\|d_{\bar{F}}^{\tilde{\rho}} - d_{\bar{F}}^\rho\| \leq \tilde{\epsilon}$, and $\|d_{\tilde{\mathcal{T}}}^{\tilde{\varphi}} - d_{\mathcal{T}^\varphi}\| \leq 2|V|C\tilde{\epsilon}$. Since $\tilde{\epsilon}$ was arbitrary small, $(\tilde{\rho}, \tilde{\varphi}, \tilde{\mathcal{T}})$ is a required rational feasible triple in Lemma 3.2.

We show Claim 3.4. Compare $\tau(\rho, \varphi, \mathcal{T})$ with $g(d_{\bar{F}}^\rho)$ (defined by (3.7)). By (the proof of) Lemma 3.1, $g(d_{\bar{F}}^\rho) \leq \tau(\rho, \varphi, \mathcal{T})$ holds. Suppose that $g(d_{\bar{F}}^\rho) < \tau(\rho, \varphi, \mathcal{T})$. This means that $\pi^{\varphi, \mathcal{T}}$ is feasible but nonoptimal to (3.5) with $d = d_{\bar{F}}^\rho$ fixed. By convexity, there is a feasible $\tilde{\pi}$ in an arbitrary open neighborhood of π with $\tau(d_{\bar{F}}^\rho, \tilde{\pi}) < \tau(\rho, \varphi, \mathcal{T})$.

Suppose $g(d_{\bar{F}}^\rho) = \tau(\rho, \varphi, \mathcal{T})$. This means that $d_{\bar{F}}^\rho$ is nonoptimal to the following convex optimization:

$$\text{Min. } g(d) \quad \text{s.t. } d: \text{ metric on } V \text{ with } d(s, t) \geq \mu(s, t) \text{ for } s, t \in S.$$

Take a sufficiently small $\epsilon' > 0$. By convexity, there exists a rational metric d' on V with $d(s, t) \geq \mu(s, t)$ for $s, t \in S$ such that $g(d') < g(d_{\bar{F}}^\rho)$ and $\|d' - d_{\bar{F}}^\rho\| \leq \epsilon'$. According to Claim 3.3, take a map $\rho' : V \rightarrow \bar{F}$ such that $\rho'(s) = p_s$, $d_{\bar{F}}^{\rho'} \leq d'$, and the length of each segment in $\mathcal{E}_{\bar{F}, \varphi}$ is rational. Then

$$\rho'(x) \in B(\rho(x), \epsilon') \quad (x \in V). \quad (3.9)$$

Indeed, by assumption (3.1), there exists $s \in S$ with $d_{\bar{F}}(\rho'(x), \rho(x)) + d_{\bar{F}}(p_s, \rho(x)) = d_{\bar{F}}(p_s, \rho'(x))$. Thus we have

$$\begin{aligned} d_{\bar{F}}(\rho'(x), \rho(x)) &= d_{\bar{F}}^{\rho'}(s, x) - d_{\bar{F}}^\rho(s, x) = \\ &= (d'(s, x) - d_{\bar{F}}^\rho(s, x)) + (d_{\bar{F}}^{\rho'}(s, x) - d'(s, x)) \leq \epsilon', \end{aligned}$$

where the last inequality follows from $\|d' - d_{\bar{F}}^{\rho}\| \leq \epsilon'$ and $d_{\bar{F}}^{\rho'} \leq d'$. Moreover, since ϵ' is small,

$$B(\rho(x), \epsilon') \cap B(\rho(y), \epsilon') = \emptyset \quad (x, y \in V : \rho(x) \neq \rho(y)). \quad (3.10)$$

For $\alpha \in [0, 1]$, define $\rho_\alpha : V \rightarrow \bar{T}$ by $\rho_\alpha(x) :=$ the (unique) point p in $[\rho(x), \rho'(x)] \subseteq B(\rho(x), \epsilon')$ with $d_{\bar{F}}(\rho(x), p) = (1-\alpha)d_{\bar{F}}(\rho(x), \rho'(x))$ and $d_{\bar{F}}(p, \rho'(x)) = \alpha d_{\bar{F}}(\rho(x), \rho'(x))$. Then, by (3.10), we can take a simple path in \bar{T} containing $[\rho(x), \rho'(x)]$ and $[\rho(y), \rho'(y)]$ for $x, y \in V$. From this we obtain

$$d_{\bar{F}}^{\rho_\alpha} = (1-\alpha)d_{\bar{F}}^{\rho} + \alpha d_{\bar{F}}^{\rho'}. \quad (3.11)$$

Take a rational optimal solution π' of (3.5) with $d = d_{\bar{F}}^{\rho'}$ fixed. Then $\tau(d_{\bar{F}}^{\rho'}, \pi') = g(d_{\bar{F}}^{\rho'}) < \tau(\rho, \varphi, \mathcal{T})$. For $\alpha \in [0, 1]$, define π_α by

$$\pi_\alpha := (1-\alpha)\pi^{\varphi, \mathcal{T}} + \alpha\pi'.$$

We are going to verify that for small rational $\alpha > 0$, $(d_{\bar{F}}^{\rho_\alpha}, \pi_\alpha)$ is a required rational solution. By $\rho_\alpha(x) \in B(\rho(x), \epsilon')$, we have $\|d_{\bar{F}}^{\rho_\alpha} - d_{\bar{F}}^{\rho}\| \leq 2\epsilon'$. Therefore we have

$$\mathcal{D}^\sigma(d_{\bar{F}}^{\rho} - a) \subseteq \mathcal{D}^\sigma(d_{\bar{F}}^{\rho_\alpha} - a) = \mathcal{D}^\sigma(d_{\bar{F}}^{\rho'} - a) \quad (\sigma \in \{-, +\}, \alpha > 0). \quad (3.12)$$

Then $(d_{\bar{F}}^{\rho_\alpha}, \pi_\alpha)$ is feasible to (3.5) since

$$\sum_X \pi_\alpha \delta_X(e) \leq (1-\alpha)|d_{\bar{F}}^{\rho} - a|(e) + \alpha|d_{\bar{F}}^{\rho'} - a|(e) = |d_{\bar{F}}^{\rho_\alpha} - a|(e) \quad (e \in E),$$

where the last equality follows from (3.11) and (3.12). Therefore $(d_{\bar{F}}^{\rho_\alpha}, \pi_\alpha)$ satisfies (2), (3) and (4) of Claim 3.4 for small $\alpha > 0$. Finally, we verify that τ is linear on α :

$$\tau(d_{\bar{F}}^{\rho_\alpha}, \pi_\alpha) = (1-\alpha)\tau(\rho, \varphi, \mathcal{T}) + \alpha\tau(d_{\bar{F}}^{\rho'}, \pi'). \quad (3.13)$$

This implies (1) in Claim 3.4 since $\tau(d_{\bar{F}}^{\rho'}, \pi') < \tau(\rho, \varphi, \mathcal{T})$. We show (3.13). By (3.12) we have $(d_{\bar{F}}^{\rho_\alpha} - a)^+(E) = (1-\alpha)(d_{\bar{F}}^{\rho} - a)^+(E) + \alpha(d_{\bar{F}}^{\rho'} - a)^+(E)$. Moreover, by (3.12) (with the same argument after Claim 3.4), $h[d_{\bar{F}}^{\rho_\alpha} - a] = h[d_{\bar{F}}^{\rho'} - a]$ for $\alpha > 0$, and $h[d_{\bar{F}}^{\rho_\alpha} - a](X_\omega) = h[d_{\bar{F}}^{\rho} - a](X_\omega)$ for $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$. Thus we have

$$\begin{aligned} \sum h[d_{\bar{F}}^{\rho_\alpha} - a](X)\pi_\alpha(X) &= (1-\alpha) \sum h[d_{\bar{F}}^{\rho} - a](X)\pi^{\varphi, \mathcal{T}}(X) + \\ &\quad + \alpha \sum h[d_{\bar{F}}^{\rho'} - a](X)\pi'(X). \end{aligned}$$

This implies (3.13). \square

3.2 Dual integrality

A feasible triple $(\rho, \varphi, \mathcal{T})$ is said to be *bipartite* if $\mathcal{T} = \bar{T}$ for some graph-theoretical tree T , $\varphi(V) \subseteq V(T)$, $\rho(V) \subseteq V(\Gamma)$, and $d_{\bar{F}}^{\rho} + d_{\mathcal{T}}^{\varphi}$ is even-valued. Consider the Cartesian product graph $\Gamma \times T$ of two trees Γ and T , which is also a bipartite graph. From the last condition, the image of (ρ, φ) belongs to one of the color classes (containing (p_s, r)); see Figure 1. The main result in this section is that the minimum of τ is attained by a bipartite triple.




Figure 1: Product of two trees

Theorem 3.5. *Suppose that μ is an even-valued tree metric realized by $(\Gamma, \{p_s\}_{s \in S}; 1)$, and edge-cost a is even-valued. For every rational feasible triple $(\rho, \varphi, \mathcal{T})$, there exists a bipartite feasible triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ such that*

- (1) $\tau(\rho^*, \varphi^*, \mathcal{T}^*) \leq \tau(\rho, \varphi, \mathcal{T})$, and
- (2) for each $x, y \in V$, if $(d_{\Gamma}^{\rho} + d_{\mathcal{T}}^{\varphi})(x, y) \leq L$ for an even integer L , then $(d_{\Gamma}^{\rho^*} + d_{\mathcal{T}^*}^{\varphi^*})(x, y) \leq L$.

In particular, the minimum of τ is attained by a bipartite feasible triple.

The second condition (2) describes a stability, which plays a crucial role in the proof of the primal integrality. Since the image of ρ belongs to $V(\Gamma)$, we have the following.

Corollary 3.6. *Under the same assumption, we have*

$$\overline{\text{val}}(\mu; G, a) = \min \max(d_{\Gamma, 1}^{\rho} - a)(E \setminus F),$$

where the maximum is taken over all inner-odd-joins F and the minimum is taken over all maps $\rho : V \rightarrow V(\Gamma)$ with $\rho(s) = p_s$ for $s \in S$.

The remainder of this subsection is devoted to the proof of Theorem 3.5, which is the most technical part of the paper. To grasp the outline of the whole proof, readers may skip the rest and proceed to Section 3.3 in the first reading.

0: Notation. Let $(\rho, \varphi, \mathcal{T})$ be a (nonbipartite) feasible triple. We may assume that

$$h[d_{\Gamma}^{\rho} - a](X_{\omega}) = 1 \quad (\omega \in \mathcal{E}_{\mathcal{T}, \varphi}). \quad (3.14)$$

Otherwise, by contracting segment ω with $h[d_{\Gamma}^{\rho} - a](X_{\omega}) = 0$, we obtain a new $(\tilde{\varphi}, \tilde{\mathcal{T}})$ with $d_{\tilde{\mathcal{T}}}^{\tilde{\varphi}} \leq d_{\mathcal{T}}^{\varphi}$. Then $(\rho, \tilde{\varphi}, \tilde{\mathcal{T}})$ is feasible, and $\tau(\rho, \tilde{\varphi}, \tilde{\mathcal{T}}) = \tau(\rho, \varphi, \mathcal{T})$. Therefore it suffices to prove Theorem 3.5 for $(\rho, \tilde{\varphi}, \tilde{\mathcal{T}})$.

By applying repeated local changes to $(\rho, \varphi, \mathcal{T})$, we try to construct a bipartite feasible triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ satisfying conditions (1) and (2). We need some notation to describe such local changes. For notational simplicity, we often denote $d_{\mathcal{T}}, d_{\bar{\Gamma}}$ by d , such as d^{ρ} and d^{ρ} , if they are distinguished in the context. Let $\{R, B\}$ be the color classes of (bipartite graph) Γ . Since μ is even-valued, we may assume $\{p_s\}_{s \in S} \subseteq B$. By rationality of $(\rho, \varphi, \mathcal{T})$, the length of each segment in $\mathcal{E}_{\bar{\Gamma}, \rho}$ and in $\mathcal{E}_{\mathcal{T}, \varphi}$ is a multiple of $\epsilon := 1/M$ for a positive integer M . An ϵ -point is a point q in \mathcal{T} such that $d_{\mathcal{T}}(q, p)$ is a multiple of ϵ for some (every) vertex p in \mathcal{T} ; the image of φ belongs to the set of ϵ -points. Similarly we can consider ϵ -points in $\bar{\Gamma}$. So we can regard \mathcal{T} (and $\bar{\Gamma}$) as a (graph-theoretical) tree on the set of ϵ -points with uniform edge-length ϵ . For an ϵ -point p in $\bar{\Gamma} \setminus V(\Gamma)$, there is a unique edge uv in Γ with $p \in [u, v]$ and $(u, v) \in R \times B$. Let $p^{\rightarrow B, \epsilon}$ denote the adjacent ϵ -point p' close to B , the unique point p' satisfying $d(p, v) = d(p, p') + \epsilon$. Similarly, let $p^{\rightarrow R, \epsilon}$ denote the unique point p' with $d(p, u) = d(p, p') + \epsilon$.

Recall the partial order \preceq (Section 2.2). For $q \in \mathcal{T}$, let $X_q, X_{q, \preceq}$, and $X_{q, \prec}$ denote the sets of nodes $x \in V$ with $q = \varphi(x)$, $q \preceq \varphi(x)$, and $q \prec \varphi(x)$, respectively. For an ϵ -point $q \in \mathcal{T}$, let $\mathcal{C} = \mathcal{C}_q$ be the set of connected components of $\mathcal{T} \setminus B^{\circ}(q, \epsilon)$. Let C_0 denote the (unique) component in \mathcal{C} containing r ; it exists exactly when $q \neq r$. By (3.14) and Lemma 2.4 (1) we have:

$$h[d_{\bar{\Gamma}}^{\rho} - a](X_{q, \preceq}) = 1, \quad (3.15)$$

$$h[d_{\bar{\Gamma}}^{\rho} - a](X_{q, \prec}) \equiv |\mathcal{C} \setminus \{C_0\}| \pmod{2} \quad (q \in \mathcal{T} \setminus \{r\}). \quad (3.16)$$

1: Critical points. Consider an ϵ -point $q \in \mathcal{T}$ with the following properties:

- (3.17) (i) $\rho(X_{q, \prec}) \subseteq V(\Gamma)$.
- (ii) For $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$, π_{ω} is an integer if ω belongs to a component in $\mathcal{C} \setminus \{C_0\}$.
- (iii) For $x, y \in X_{q, \prec}$, $(d^{\rho} + d^{\varphi})(x, y)$ is an even integer if $\varphi(x)$ and $\varphi(y)$ belong to the same component in $\mathcal{C} \setminus \{C_0\}$.

This means that the restriction of $(\rho, \varphi, \mathcal{T})$ to (the preimage of) each component in $\mathcal{C} \setminus \{C_0\}$ is bipartite. Therefore we can draw a *grid* on $C \times \bar{\Gamma}$ (as in Figure 2) so that its vertices are points having *even* distance-sum $d_{\bar{\Gamma}} + d_{\mathcal{T}}$ from some (every) image of (ρ, φ) in $C \times \bar{\Gamma}$. We try to extend these *partial* grids to the *whole* grid; the subsequent argument is based on this intuition. A *critical point* is a minimal ϵ -point having the property (3.17). We will make local changes on $(\rho, \varphi, \mathcal{T})$ at critical points, and move critical points toward r .

Let $q \in \mathcal{T}$ be a critical point. Fix $b \in B$. For a component $C \in \mathcal{C} \setminus \{C_0\}$, take a node x with $\varphi(x) \in C$ (it exists by (3.14)). By condition (3.17), we see that $d(q, \varphi(x)) - \lfloor d(q, \varphi(x)) \rfloor$ and the parity of $d(b, \rho(x)) + \lfloor d(q, \varphi(x)) \rfloor$ are determined independent of the choice of x and b . We denote them by Δ_C and D_C , respectively. Partition $\mathcal{C} \setminus \{C_0\}$ according to Δ_C, D_C as:

$$\begin{aligned} \mathcal{C}^B &:= \{C \mid \Delta_C \neq 0, D_C = \text{even}\}, & \mathcal{C}^R &:= \{C \mid \Delta_C \neq 0, D_C = \text{odd}\}, \\ \mathcal{I}^B &:= \{C \mid \Delta_C = 0, D_C = \text{even}\}, & \mathcal{I}^R &:= \{C \mid \Delta_C = 0, D_C = \text{odd}\}. \end{aligned}$$

Figure 2 illustrates a portion of $C \times \bar{\Gamma}$ for $C \in \mathcal{C} \setminus \{C_0\}$.



Figure 2: Drawing grids in $C \times \bar{\Gamma}$ for (B) $C \in \mathcal{C}^B \cup \mathcal{I}^B$ and (R) $C \in \mathcal{C}^R \cup \mathcal{I}^R$ 

Figure 3: Tree modification I

Let X^B be the set of nodes $x \in X_{q, \geq}$ such that $d(b, \rho(x)) + d(q, \varphi(x))$ is even. X^B consists of nodes $x \in X_q$ with $\rho(x) \in B$ and nodes $x \in X_{q, >}$ with $\varphi(x) \in C \in \mathcal{I}^B$. Similarly, let X^R be the set of nodes $x \in X_{q, \geq}$ such that $d(b, \rho(x)) + d(q, \varphi(x))$ is odd. For $C \in \mathcal{C}$ let q_C denote the (unique) point in C meeting $B(q, \epsilon)$.

2: Local changes at $q \neq r$. Take a critical point q . We first consider the case $q \neq r$. Suppose that $X^B \neq \emptyset$ and $h[d^p - a](X^B) = 0$. This implies $h[d^p - a](X_{q, \geq} \setminus X^B) = 1$ by Lemma 2.4 (1) and (3.15). We construct modification (φ', \mathcal{T}') of (φ, \mathcal{T}) as follows. First let $\varphi'(x) := \varphi(x)$ for $x \in V \setminus X_q$. Next delete open ball $B^\circ(q, \epsilon)$ from \mathcal{T} . Let \mathcal{C} be the set of the resulting connected components. As above, partition \mathcal{C} into $\{C_0\} \cup \mathcal{C}^B \cup \mathcal{C}^R \cup \mathcal{I}^B \cup \mathcal{I}^R$. For each $C \in \mathcal{I}^B$, join q_C and q_{C_0} by a segment of length ϵ . Add a new point \bar{q} , and join it with q_{C_0} by a segment of length ϵ . For each $C \in \mathcal{C} \setminus \mathcal{I}^B$, join \bar{q} and q_C by a segment of length ϵ . Now we obtain a new tree \mathcal{T}' and obtain a map $\varphi' : V \setminus X_q \rightarrow \mathcal{T}'$. We extend φ' to $V \rightarrow \mathcal{T}'$ by defining $\varphi'(x) := q_{C_0}$ for $x \in X^B \cap X_q$ and $\varphi'(x) := \bar{q}$ for $x \in X_q \setminus X^B$. See the middle of Figure 3. We claim

Claim 3.7. $(\rho, \varphi', \mathcal{T}')$ is feasible, and $\tau(\rho, \varphi', \mathcal{T}') = \tau(\rho, \varphi, \mathcal{T})$.

Proof. To see the feasibility (3.2) (2), take an edge $e = xy \in E$. It suffices to consider the case where $x \in X^B$ and $y \in X_{q, \geq} \setminus X^B$; for the other cases we have $d^{\varphi'}(e) \leq d^\varphi(e)$, and thus (3.2) (2) is preserved. Then $d^{\varphi'}(e) = d^\varphi(e) + \epsilon$.




Figure 4: Tree modification II

So it suffices to show $d^\varphi(e) < |d^\rho - a|(e)$. This follows from the fact that $a(e)$ is even and $(d^\rho + d^\varphi)(e)$ is not even. The latter part of Claim 3.7 follows from: $h \circ (\varphi', \mathcal{T}') - h \circ (\varphi, \mathcal{T}) = \epsilon (h(X_{q, \leq} \setminus X^B) - h(X_{q, \leq})) = 0$, where we denote $h[d^\rho - a]$ by h , and we use Lemma 2.4 (1) with $h(X^B) = 0$ and $h(X_{q, \leq}) = 1$ (3.15). \square

Let $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho, \varphi', \mathcal{T}')$. Then X^B becomes empty at q . Suppose that $X^B \neq \emptyset$ and $h[d^\rho - a](X^B) = 1$. This implies $h[d^\rho - a](X_{p, \leq} \setminus X^B) = 0$. Therefore we can apply the above-described process with changing the roles of X^B and $X_{p, \leq} \setminus X^B$. See the right of Figure 3. Then we get a feasible triple $(\rho, \varphi', \mathcal{T}')$. Let $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho, \varphi', \mathcal{T}')$. Suppose $X^R \neq \emptyset$. Then apply the above-described procedure by changing roles of B and R . We remark that (3.14) is preserved.

So consider the case of $X^B \cup X^R = \emptyset$, which implies $\mathcal{I}^B \cup \mathcal{I}^R = \emptyset$. Here we construct two modifications $(\rho', \varphi', \mathcal{T}')$ and $(\rho'', \varphi'', \mathcal{T}'')$ as follows. First let $(\rho', \varphi')(x) := (\rho, \varphi)(x)$ for $x \in V \setminus X_q$. Next delete open ball $B^\circ(q, \epsilon)$ from \mathcal{T} . Let $\mathcal{C} = \{C_0\} \cup \mathcal{C}^B \cup \mathcal{C}^R$ be the set of resulting connected components. For each $C \in \mathcal{C}^R$, join q_C and q_{C_0} by a segment of length 2ϵ . For each $C \in \mathcal{C}^B$, identify (glue) q_C and q_{C_0} . For each $x \in X_q$, let $\varphi'(x) := q_{C_0}$ and let $\rho'(x) := (\rho(x))^{\rightarrow B, \epsilon}$ (well-defined by $X^B \cup X^R = \emptyset$). Let $(\rho', \varphi', \mathcal{T}')$ be the resulting triple. See Figure 4 for the construction of \mathcal{T}' . Namely \mathcal{T}' is also obtained from \mathcal{T} by contracting by ϵ each segment connecting q and $C \in \{C_0\} \cup \mathcal{C}^B$ and extending by ϵ each segment connecting q and $C \in \mathcal{C}^R$. The second triple $(\rho'', \varphi'', \mathcal{T}'')$ is obtained by changing roles of B and R .

Claim 3.8. (1) Both $(\rho', \varphi', \mathcal{T}')$ and $(\rho'', \varphi'', \mathcal{T}'')$ are feasible.

(2) $\mathcal{D}^\sigma(d^{\rho'} - a) \subseteq \mathcal{D}^\sigma(d^\rho - a)$ and $\mathcal{D}^\sigma(d^{\rho''} - a) \subseteq \mathcal{D}^\sigma(d^\rho - a)$ for $\sigma \in \{-, +\}$.

(3) Either $\tau(\rho', \varphi', \mathcal{T}') \leq \tau(\rho, \varphi, \mathcal{T})$ or $\tau(\rho'', \varphi'', \mathcal{T}'') \leq \tau(\rho, \varphi, \mathcal{T})$ holds.

Let $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho', \varphi', \mathcal{T}')$ if $\tau(\rho', \varphi', \mathcal{T}') \leq \tau(\rho, \varphi, \mathcal{T})$, and let $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho'', \varphi'', \mathcal{T}'')$ otherwise. Repeat this process. Then critical points move toward r . After finitely many steps, r becomes critical; go to paragraph 3 after the proof of Claim 3.8.

Proof. (1). It suffices to show the feasibility of $(\rho', \varphi', \mathcal{T}')$. Take an edge $e = xy \in E$ with $(\rho, \varphi)(x) \neq (\rho, \varphi)(y)$. We verify condition (*) $d^{\rho'}(e) \leq |d^{\rho'}(e) - a(e)|$ for the cases (i) $\varphi(x) = q \prec \varphi(y) \in C \in \mathcal{C}^R$, and (ii) $\varphi(x) \in C \in \mathcal{C}^R$, $\varphi(y) \in C' \in \mathcal{C}^R$, and $C \neq C'$. For the other cases, $d^{\rho'}(e) < d^\rho(e)$, or $d^{\rho'}(e) = d^\rho(e)$ and $d^{\rho'}(e) = d^\rho(e)$; the condition (*) is easily verified.

Case (i): $\varphi(x) = q \prec \varphi(y) \in C \in \mathcal{C}^R$. In this case, $d^{\varphi'}(e) = d^{\varphi}(e) + \epsilon$. We are going to show that (*1) $|d^{\rho'} - a|(e) = |d^{\rho} - a|(e) + \epsilon$ or (*2) $|d^{\rho} - a|(e) - d^{\varphi}(e) \geq 2\epsilon$, which implies the feasibility (*). Take $uv \in E(\Gamma)$ with $\rho(x) \in (u, v)$. We may assume $d^{\rho}(e) = d(\rho(x), u) + d(u, \rho(y))$ (since $\rho(y) \in V(\Gamma)$). By the tree structure, we have $u \in B \Leftrightarrow d^{\rho'}(e) = d^{\rho}(e) - \epsilon$, and $u \in R \Leftrightarrow d^{\rho'}(e) = d^{\rho}(e) + \epsilon$. If $u \in B$ and $d^{\rho}(e) < a(e)$ then (*1) holds. Also, if $u \in R$ and $d^{\rho}(e) > a(e)$ then (*1) holds. So suppose that $u \in B$ and $d^{\rho}(e) > a(e)$. Let $t := d^{\varphi}(e) - \lfloor d(\varphi(y), q) \rfloor$. Then we obtain

$$|d^{\rho} - a|(e) - d^{\varphi}(e) = d(u, \rho(y)) + d(\rho(x), u) - a(e) - \lfloor d(\varphi(y), q) \rfloor - t.$$

By $\rho(y) \in C \in \mathcal{C}^R$, $u \in B$, and $a(e) \in 2\mathbf{Z}$, integer $N := d(u, \rho(y)) - \lfloor d(\varphi(y), q) \rfloor - a(e)$ is odd. Furthermore, both $d(\rho(x), u)$ and t belong to $[\epsilon, 1 - \epsilon]$. Hence we have

$$N - 1 + 2\epsilon \leq |d^{\rho} - a|(e) - d^{\varphi}(e) \leq N + 1 - 2\epsilon.$$

Here $N \leq -1$ is impossible; otherwise $|d^{\rho} - a|(e) - d^{\varphi}(e) < 0$, contradicting the feasibility of $(\rho, \varphi, \mathcal{T})$. Thus $N \geq 1$, implying (*2). The analysis for the case of $u \in R$ and $d^{\rho}(e) < a(e)$ is similar.

Case (ii): $\varphi(x) \in C \in \mathcal{C}^R$, $\varphi(y) \in C' \in \mathcal{C}^R$, and $C \neq C'$. In this case, $d^{\rho'}(e) = d^{\rho}(e)$ and $d^{\varphi'}(e) = d^{\varphi}(e) + 2\epsilon$. Here $d^{\varphi}(e) = d(\varphi(x), q) + d(\varphi(y), q)$. Let $t := d(\varphi(x), q) - \lfloor d(\varphi(x), q) \rfloor$ and $t' := d(\varphi(y), q) - \lfloor d(\varphi(y), q) \rfloor$. Then we have

$$|d^{\rho} - a|(e) - d^{\varphi}(e) = |d^{\rho} - a|(e) - \lfloor d(\varphi(x), q) \rfloor - \lfloor d(\varphi(y), q) \rfloor - t - t'.$$

The integer $N := |d^{\rho} - a|(e) - \lfloor d(\varphi(x), q) \rfloor - \lfloor d(\varphi(y), q) \rfloor$ is even since $C, C' \in \mathcal{C}^R$ and $|d^{\rho} - a|(e) \equiv d(b, \rho(x)) + d(b, \rho(y)) \pmod{2}$. By $t, t' \in [\epsilon, 1 - \epsilon]$, we have

$$N - 2 + 2\epsilon \leq |d^{\rho} - a|(e) - d^{\varphi}(e) \leq N - 2\epsilon.$$

From this, $N \geq 2$ is necessary, and we have $|d^{\rho} - a|(e) - d^{\varphi}(e) \geq 2\epsilon$, implying (*).

(2). Take an edge $e = xy \in E$. We show that (**) neither $(d^{\rho'} - a)(e) < 0 \leq (d^{\rho} - a)(e)$ nor $(d^{\rho'} - a)(e) > 0 \geq (d^{\rho} - a)(e)$ occurs. We may consider the case $\rho(x) = q$. Suppose $\rho(y) \neq q$. Then $|d^{\rho} - a|(e) \geq d^{\varphi}(e) \geq \epsilon$, and $d^{\rho'}(e) - d^{\rho}(e) \in \{\epsilon, -\epsilon\}$, implying (**). So suppose $\varphi(x) = \varphi(y) = q$. Then $d^{\varphi'}(e) = d^{\varphi}(e) = 0$, and $d^{\rho'}(e) - d^{\rho}(e) = \{2\epsilon, 0, -2\epsilon\}$. We may consider the case where $\rho(x) \in (u, p)$ and $\rho(y) \in (u', p')$ for different edges $e = up, e' = u'p' \in E(\Gamma)$ (otherwise $d^{\rho'}(e) - d^{\rho}(e) = 0$). We can assume that $d^{\rho}(x, y) = d(\rho(x), p) + d(p, p') + d(p', \rho(y))$. If p and p' belong to different color classes, then $\rho(x)$ and $\rho(y)$ move toward the same direction in the path from $\rho(x)$ to $\rho(y)$, and hence we have $d^{\rho'}(e) = d^{\rho}(e)$ implying (**). Suppose that p and p' belong to the same color class. Then $N := d(p, p')$ is even. By $d(\rho(x), p), d(p', \rho(y)) \in [\epsilon, 1 - \epsilon]$ we have

$$N - a(e) + 2\epsilon \leq (d^{\rho} - a)(e) \leq N - a(e) + 2 - 2\epsilon.$$

Since $a(e)$ is even, we have $|d^{\rho} - a|(e) \geq 2\epsilon$ implying (**).

(3). We first compare $(d^{\rho'} - a)^+(E)$ with $(d^{\rho} - a)^+(E)$. For a node $x \in V$ with $\rho(x) \notin V(\Gamma)$, define disjoint subsets $\delta^B x, \delta^R x$ of δx as follows. Take edge $uv \in E(\Gamma)$ with $\rho(x) \in [u, v]$. Suppose $u \in B$ and $v \in R$. Let $\delta^B x$ (resp. $\delta^R x$) be

the set of edges $e = xy \in \delta x$ such that $\rho(y)$ belongs to the connected component of $\bar{\Gamma} \setminus \{\rho(x)\}$ containing u (resp. v). Then we have

$$\begin{aligned} & (d^{\rho'} - a)^+(E) - (d^{\rho} - a)^+(E) = (d^{\rho'} - d^{\rho})(E \setminus \mathcal{D}^-(d^{\rho} - a)) \quad (3.18) \\ & = \epsilon \sum_{x \in X_q} (|\delta^R x \setminus \mathcal{D}^-(d^{\rho} - a)| - |\delta^B x \setminus \mathcal{D}^-(d^{\rho} - a)|) \\ & = (d^{\rho} - a)^+(E) - (d^{\rho''} - a)^+(E). \end{aligned}$$

Here we use (2) for the first equality. To see the second equality, consider an edge $e \in E \setminus \mathcal{D}^-(d^{\rho} - a)$ joining X_q ; the other edges do not contribute. Further we may assume $d^{\rho}(e) > 0$; otherwise $d^{\rho}(e) = d^{\rho'}(e) = 0$ (by $a(e) \geq 0$) and e does not contribute. Suppose $e = xy$ and $x \in X_q$. Then $e \in \delta^R x \cup \delta^B x$. By the case-by-case analysis, we see

$$d^{\rho'}(e) - d^{\rho}(e) = \begin{cases} -\epsilon & \text{if } e \in \delta^B x, y \notin X_q, \\ +\epsilon & \text{if } e \in \delta^R x, y \notin X_q, \\ -2\epsilon & \text{if } e \in \delta^B x \cap \delta^B y, y \in X_q \\ +2\epsilon & \text{if } e \in \delta^R x \cap \delta^R y, y \in X_q \\ 0 & \text{otherwise.} \end{cases} \quad (3.19)$$

By a simple counting argument, we see the second equality. The third equality is obtained by changing the roles of R and B .

Next we compare $h[d^{\rho'} - a] \circ (\varphi', \mathcal{T}')$ with $h[d^{\rho} - a] \circ (\varphi, \mathcal{T})$. Here \mathcal{T}' is obtained from \mathcal{T} by contracting and extending segments in $\mathcal{E}_{\mathcal{T}, \varphi}$; so we can naturally regard $\mathcal{E}_{\mathcal{T}', \varphi'}$ as $\mathcal{E}_{\mathcal{T}', \varphi} \subseteq \mathcal{E}_{\mathcal{T}, \varphi}$. Moreover it holds that

$$h[d^{\rho'} - a](X_{\omega}) = h[d^{\rho} - a](X_{\omega}) = 1 \quad (\omega \in \mathcal{E}_{\mathcal{T}', \varphi'}). \quad (3.20)$$

Indeed, for $\omega \in \mathcal{E}_{\mathcal{T}', \varphi'}$, we have $|d^{\rho'} - a|(e) \geq d^{\rho'}(e) > 0$ ($e \in \delta X_{\omega}$), and by (2) $d^{\rho'} - a$ and $d^{\rho} - a$ must have the same sign pattern on δX_{ω} . This implies (3.20).

\mathcal{T}' is obtained from \mathcal{T} by decreasing the length of ω by ϵ if ω connects q and a component in $\{C_0\} \cup \mathcal{C}^B$ and by increasing the length of ω by ϵ if ω connects q and a component in \mathcal{C}^R . By this fact together with (3.20), we have

$$h[d^{\rho'} - a] \circ (\varphi', \mathcal{T}') - h[d^{\rho} - a] \circ (\varphi, \mathcal{T}) = \epsilon |\mathcal{C}^R| - \epsilon |\mathcal{C}^B| - \epsilon. \quad (3.21)$$

Similarly we have

$$h[d^{\rho''} - a] \circ (\varphi'', \mathcal{T}'') - h[d^{\rho} - a] \circ (\varphi, \mathcal{T}) = \epsilon |\mathcal{C}^B| - \epsilon |\mathcal{C}^R| - \epsilon. \quad (3.22)$$

Let $\Delta' := \{\tau(\rho', \varphi', \mathcal{T}') - \tau(\rho, \varphi, \mathcal{T})\}/\epsilon$ and $\Delta'' := \{\tau(\rho'', \varphi'', \mathcal{T}'') - \tau(\rho, \varphi, \mathcal{T})\}/\epsilon$. By (3.18), (3.21), and (3.22), we have

$$\Delta'' = -\Delta' + 2.$$

Therefore it suffices to show that Δ' is an even integer, which implies (3) either $\Delta' \leq 0$ or $\Delta'' \leq 0$. By (3.18) and (3.21), we have

$$\begin{aligned} \Delta' & \equiv \sum_{x \in X_q} (|\delta^R x \setminus \mathcal{D}^-(d^{\rho} - a)| + |\delta^B x \setminus \mathcal{D}^-(d^{\rho} - a)|) + (|\mathcal{C}^B| + |\mathcal{C}^R|) + 1 \\ & \equiv \sum_{x \in X_q} |\delta x \setminus \mathcal{D}^-(d^{\rho} - a)| + h[d^{\rho} - a](X_{q, <}) + 1 \\ & \equiv \sum_{x \in X_q} h[d^{\rho} - a](x) + h[d^{\rho} - a](X_{q, <}) + 1 \equiv h[d^{\rho} - a](X_{q, \leq}) + 1 \equiv 0 \pmod{2}. \end{aligned}$$

In the second equality, we use the observation that for $e = xy \in E \setminus \mathcal{D}^-(d^p - a)$ if $x \in X_q$ and $e \notin \delta^B x \cup \delta^R x$, then $d^p(x, y) = d^\varphi(x, y) = 0$, and $y \in X_q$ and $e \notin \delta^B y \cup \delta^R y$. Also we use (3.15), (3.16), and Lemma 2.4 (1). \square

3: Local changes at r . Suppose that r is the unique critical point. Let $q := r$. The argument is essentially the same (or simpler). Suppose $X_{q, \leq} \setminus (X^B \cup X^R) \neq \emptyset$. As above, we construct $(\rho', \varphi', \mathcal{T}')$ and $(\rho'', \varphi'', \mathcal{T}'')$ in the following way. Define $\rho' : V \rightarrow \bar{\Gamma}$ by $\rho'(x) := (\rho(x))^{-B, \epsilon}$ for $x \in X_q \setminus (X^B \cup X^R)$, and $\rho'(x) := \rho(x)$ for $x \in X^B \cup X^R$. Also define $\varphi'(x) := \varphi(x)$ for $x \in V \setminus X_q$. Next delete open ball $B^\circ(q, \epsilon)$ from \mathcal{T} . Let $\mathcal{C} = \mathcal{C}^B \cup \mathcal{C}^R \cup \mathcal{I}^B \cup \mathcal{I}^R$ be the set of the resulting connected component in \mathcal{T} . Add a new root r . For all $C \in \mathcal{I}^B \cup \mathcal{I}^R$, join q_C and r by a segment of length ϵ . For each $C \in \mathcal{C}^B$, join q_C and r by a segment of length 2ϵ . For each $C \in \mathcal{C}^R$, identify (glue) q_C and r . Define $\varphi'(x) = r$ for $x \in X_q$. Then we obtain a triple $(\rho', \varphi', \mathcal{T}')$. The second triple $(\rho'', \varphi'', \mathcal{T}'')$ is obtained by changing the roles of B and R . By a similar analysis in Claim 3.8, both $(\rho', \varphi', \mathcal{T}')$ and $(\rho'', \varphi'', \mathcal{T}'')$ are feasible, and $\tau(\rho', \varphi', \mathcal{T}') - \tau(\rho, \varphi, \mathcal{T}) = -\tau(\rho'', \varphi'', \mathcal{T}'') + \tau(\rho, \varphi, \mathcal{T})$ (in (3.18) and (3.19), replace X_q by $X_q \setminus (X^B \cup X^R)$ and in (3.21) and (3.22), delete $-\epsilon$ in RHS since C_0 does not arise). Let $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho', \varphi', \mathcal{T}')$ if $\tau(\rho', \varphi', \mathcal{T}') \leq \tau(\rho, \varphi, \mathcal{T})$ and let $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho'', \varphi'', \mathcal{T}'')$ otherwise. Increase ϵ to its multiple until $X^B \cup X^R$ increases. Repeat it until $X_{r, \leq} = X^B \cup X^R$. Also these changes keep (2) in Claim 3.8.

If $X^R = \emptyset$, then the current $(\rho, \varphi, \mathcal{T})$ is bipartite, as required. So suppose $X^R \neq \emptyset$. Delete open ball $B^\circ(r, \epsilon)$ from \mathcal{T} . Then $\mathcal{C} = \mathcal{I}^R \cup \mathcal{I}^B$. Consider a segment of length 1 with vertices r, q ; r is a new root. For each $C \in \mathcal{C}^B$, join q_C and r by a segment of length ϵ . For each $C \in \mathcal{C}^R$, join q_C and q by a segment of length ϵ . For $x \in X_q$, define $\varphi'(x) := r$ if $x \in X^B$, and $\varphi'(x) := q$ if $x \in X^R$. The resulting $(\rho, \varphi', \mathcal{T}')$ is feasible. Indeed, if e join X^B and X^R , then $d^\varphi(e)$ and $|d^p(e) - a(e)|$ have different parity, and this implies $d^\varphi(e) + 1 \leq |d^p(e) - a(e)|$. From this we see the feasibility. Now $(\rho, \varphi', \mathcal{T}')$ is bipartite, and $\tau(\rho, \varphi', \mathcal{T}') = \tau(\rho, \varphi, \mathcal{T}) - h[d^p - a](X^B) \leq \tau(\rho, \varphi, \mathcal{T})$. Hence (1) of Theorem 3.5 is proved.

4: Stability. Finally we verify that the resulting $(\rho^*, \varphi^*, \mathcal{T}^*)$ satisfies the condition (2) in Theorem 3.5. It suffices to verify that for each change $(\rho, \varphi, \mathcal{T}) \leftarrow (\rho', \varphi', \mathcal{T}')$ above, $(d^p + d^\varphi)(x, y) \leq L$ implies $(d^{\rho'} + d^{\varphi'})(x, y) \leq L$ for an even integer L . This has already been done. Add an edge e joining x and y with *even* cost $a(e) = L$. Then $(\rho, \varphi, \mathcal{T})$ is feasible to the new graph, and then so does $(\rho', \varphi', \mathcal{T}')$ with property (2) in Claim 3.8. In any change, we have $(d^{\rho'} + d^{\varphi'})(x, y) \leq a(e) = L$. Thus the proof of Theorem 3.5 is complete. \square

3.3 Primal integrality

Here we prove the existence of an integral optimal solution in (2.2):

Theorem 3.9. *Suppose that μ is a tree metric. Then there exists an integral optimal solution in (2.2). In particular, $\text{val}(\mu; G, a) = \overline{\text{val}}(\mu; G, a)$.*

Together with Corollary 3.6, this establishes the min-max formula (1.2). The proof of Theorem 3.9 is based on an inductive approach by edge-deletion. Take an

edge $e \in E$, consider $G - e$. Then we have

$$\overline{\text{val}}(\mu; G - e, a) \leq \overline{\text{val}}(\mu; G, a). \quad (3.23)$$

To see this, take any optimal solution (ξ, f) in (2.2) for $G - e$. Extend ξ to $\bar{\xi} : E \rightarrow \mathbf{R}_+$ by defining $\bar{\xi}(e) := 1$. The resulting $(\bar{\xi}, f)$ is feasible to (2.2) for G . In particular, if (3.23) holds with equality and there exists an integral optimal solution in (2.2) for $G - e$, then there also exists an integral optimal solution in (2.2) for G .

An edge $e \in E$ is called *deletable* if $\overline{\text{val}}(\mu; G - e, a) = \overline{\text{val}}(\mu; G, a)$. Our goal is to prove the following simple criterion of deletable edges.

Proposition 3.10. *Let (ξ, f) be an optimal solution in (2.2). For an edge $e \in E$, if $\xi(e) > 0$, then e is deletable.*

We first prove Theorem 3.9 assuming Proposition 3.10 which we next prove.

Proof of Theorem 3.9 assuming Proposition 3.10. We use the induction on the number of edges. Take any optimal solution (ξ, f) in (2.2). Suppose $\xi = 0$. Then $\overline{\text{val}}(\mu; G, a) = \overline{\text{val}}(\mu; G, 0) - a(E)$. Moreover G is necessarily inner Eulerian. By the laminar locking theorem (Lemma 2.1), we can take an edge-disjoint set \mathcal{P} of S -paths with $\sum_{P \in \mathcal{P}} \mu(s_P, t_P) = \overline{\text{val}}(\mu; G, 0)$. Then \mathcal{P} attains the maximum since $\sum_{P \in \mathcal{P}} \mu(s_P, t_P) - a(P) \leq \overline{\text{val}}(\mu; G, a) \leq \overline{\text{val}}(\mu; G, 0) - a(E) \leq \sum_{P \in \mathcal{P}} \mu(s_P, t_P) - a(P)$.

Suppose that there is an edge $e \in E$ with $\xi(e) > 0$. By Proposition 3.10, e is deletable. By induction together with the construction above, there exists an integral optimal solution. \square

Proof of Proposition 3.10. We need three lemmas below, the first is about the parity, the second is the complementary slackness condition, and the third is a discrete version of Stability Lemma (Lemma 3.2). Since we often compare parameters of $G - e$ with that of G , we denote τ^{G-e} and h^{G-e} by τ^e and h^e , respectively. For maps $\rho : V \rightarrow \bar{F}$ and $\varphi : V \rightarrow \mathcal{T}$, we denote $d_{\bar{F}}^\rho$ and $d_{\mathcal{T}}^\varphi$ by d^ρ and d^φ , respectively, and we also denote $h[d^\rho - a]$ by h . We note:

$$(3.24) \quad (1) \quad h^e(X) \neq h(X) \text{ if and only if } e \in \delta X \setminus \mathcal{D}^-(d^\rho - a).$$

$$(2) \quad h^e \circ (\varphi, \mathcal{T}) = h \circ (\varphi, \mathcal{T}) + \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi} : e \in \delta X_\omega} \pi_\omega(h^e(X_\omega) - h(X_\omega)).$$

This follows from the definition of h .

Lemma 3.11. *If a feasible triple $(\rho, \varphi, \mathcal{T})$ is bipartite, then $\tau(\rho, \varphi, \mathcal{T})$ is an even integer.*

Proof. We use the induction on the number of edges; if $E = \emptyset$ then the statement is obvious. Take an edge $e = xy \in E$. $(\rho, \varphi, \mathcal{T})$ is feasible to $G - e$. If $e \in \mathcal{D}^-(d^\rho - a)$ then $\tau^e(\rho, \varphi, \mathcal{T}) = \tau(\rho, \varphi, \mathcal{T})$ by (3.24). Suppose $e \notin \mathcal{D}^-(d^\rho - a)$. By (3.24) (1), the second term of RHS in (3.24) (2) is equal to $d^\varphi(e)$ modulo 2; recall (2.3) in

Section 2.2. Also $(d^\rho - a)^+(E) - (d^\rho - a)^+(E \setminus e) = (d^\rho - a)^+(e) \equiv d^\rho(e) \pmod{2}$. Since $d^\rho(e) \equiv d^\varphi(e) \pmod{2}$ by bipartiteness, we have $\tau^e(\rho, \varphi, \mathcal{T}) \equiv \tau(\rho, \varphi, \mathcal{T}) \pmod{2}$, as required. \square

Lemma 3.12. *Let (ξ, f) be a feasible solution in (2.2) and $(\rho, \varphi, \mathcal{T})$ a feasible triple. Then (ξ, f) and $(\rho, \varphi, \mathcal{T})$ are both optimal if and only if the following conditions hold:*

(a) For each $e \in E$ with $d^\rho(e) > 0$ we have $\xi(e) + \sum_{P \in \Pi, e \in P} f(P) = 1$.

(b) For each $e \in E$ with $d^\varphi(e) < |d^\rho - a|(e)$ we have

$$\xi(e) = \begin{cases} 1 & \text{if } e \in \mathcal{D}^-(d^\rho - a), \\ 0 & \text{otherwise.} \end{cases}$$

(c) For each $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$, we have

$$\xi(\delta X_\omega \setminus \mathcal{D}^-(d^\rho - a)) + (1 - \xi)(\delta X_\omega \cap \mathcal{D}^-(d^\rho - a)) = h[d^\rho - a](X_\omega).$$

(d) For each $P \in \Pi$ with $f(P) > 0$ we have $d^\rho(P) = \mu(s_P, t_P)$.

Proof. By direct calculations, we obtain

$$\begin{aligned} & (d^\rho - a)^+(E) - h[d^\rho - a] \circ (\varphi, \mathcal{T}) - \sum_{P \in \Pi} \mu(s_P, t_P) f(P) + a(E) - \sum_{e \in E} a(e) \xi(e) \\ &= \sum_{e \in E} d^\rho(e) (1 - \xi(e) - f(e)) + \sum_{P \in \Pi} f(P) (d^\rho(P) - \mu(s_P, t_P)) \\ &+ \sum_{e \in E \setminus D^-} \xi(e) (|d^\rho - a| - d^\rho)(e) + \sum_{e \in D^-} (1 - \xi(e)) (|d^\rho - a| - d^\rho)(e) \\ &+ \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi}} \pi_\omega [\xi(\delta X_\omega \setminus D^-) + (1 - \xi)(\delta X_\omega \cap D^-) - h[d^\rho - a](X_\omega)], \end{aligned} \quad (3.25)$$

where $f(e) := \sum_{P \in \Pi, e \in P} f(P)$ and $D^- := \mathcal{D}^-(d^\rho - a)$. To see this, use the relations:

$$\sum_{P \in \Pi} f(P) d^\rho(P) = \sum_{e \in E} f(e) d^\rho(e),$$

$$\sum_{e \in E \setminus D^-} \xi(e) d^\rho(e) + \sum_{e \in D^-} (1 - \xi(e)) d^\rho(e) = \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi}} \pi_\omega [\xi(\delta X_\omega \setminus D^-) + (1 - \xi)(\delta X_\omega \cap D^-)],$$

and

$$\begin{aligned} & (d^\rho - a)^+(E) + a(E) - \sum_{e \in E} a(e) \xi(e) \\ &= \sum_{e \in E} d^\rho(e) (1 - \xi(e)) + \sum_{e \in E \setminus D^-} \xi(e) (d^\rho - a)(e) + \sum_{e \in D^-} (1 - \xi(e)) (a - d^\rho)(e). \end{aligned}$$

By feasibility, all terms in RHS of (3.25) are nonnegative; see also (2.6). Therefore $(\rho, \varphi, \mathcal{T})$ and (ξ, f) are both optimal if and only if the all terms are zero, which is equivalent to the case where (a), (b), (c), and (d) hold. \square

Lemma 3.13 (Discrete Stability Lemma). *If a bipartite feasible triple $(\rho, \varphi, \mathcal{T})$ is not optimal, then there exists a bipartite feasible triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ such that $\tau(\rho^*, \varphi^*, \mathcal{T}^*) < \tau(\rho, \varphi, \mathcal{T})$, and $(d^{\rho^*} + d^{\varphi^*})(x, y) \leq (d^\rho + d^\varphi)(x, y) + 2$ for $x, y \in V$.*

Proof. By contracting $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$ with $h(X_\omega) = 0$, we get a nonoptimal feasible triple $(\rho, \tilde{\varphi}, \tilde{\mathcal{T}})$ with $d^{\tilde{\varphi}} \leq d^\varphi$, $\tau(\rho, \tilde{\varphi}, \tilde{\mathcal{T}}) = \tau(\rho, \varphi, \mathcal{T})$, and $h(X_\omega) = 1$ ($\omega \in \mathcal{E}_{\tilde{\mathcal{T}}, \tilde{\varphi}}$). By Lemma 3.2, there is a rational feasible triple $(\rho', \varphi', \mathcal{T}')$ with $\tau(\rho', \varphi', \mathcal{T}') < \tau(\rho, \tilde{\varphi}, \tilde{\mathcal{T}})$ such that $(\rho', \varphi', \mathcal{T}')$ is sufficiently close to $(\rho, \tilde{\varphi}, \tilde{\mathcal{T}})$. For $x, y \in V$, let $L_{x,y}$ denote the smallest even integer not less than $(d^{\rho'} + d^{\varphi'})(x, y)$. Then $L_{x,y} \leq (d^\rho + d^{\tilde{\varphi}})(x, y) + 2 \leq (d^\rho + d^\varphi)(x, y) + 2$. By Theorem 3.5, there is a bipartite feasible triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ such that $\tau(\rho^*, \varphi^*, \mathcal{T}^*) \leq \tau(\rho', \varphi', \mathcal{T}')$ and $(d^{\rho^*} + d^{\varphi^*})(x, y) \leq L_{x,y} \leq (d^\rho + d^\varphi)(x, y) + 2$. \square

Now we are ready to prove Proposition 3.10. Take an optimal solution (ξ, f) in (2.2) and take a bipartite optimal triple $(\rho, \varphi, \mathcal{T})$. Let $e = xy \in E$ be an edge with $\xi(e) > 0$. Here we can assume

$$d^\rho(e) = d^\varphi(e) = 0 = a(e). \quad (3.26)$$

Indeed, replace e by a series of two edges xz and zy , and define $a(xz) := 0$ and $a(zy) := a(e)$. Then the problem (2.2) does not change; from any feasible solution (ξ, f) in the original instance we get a feasible solution in the new one with the same objective value by setting $\xi(xz) = \xi(zy) := \xi(e)$, and the converse is also possible by contracting edge xz . In particular e is deletable in the original graph if and only if xz is deletable in the new graph. Also we can extend (ρ, φ) by defining $(\rho, \varphi)(z) := (\rho, \varphi)(x)$, which is optimal to the new instance (since it is feasible and has the same objective value).

Suppose indirectly that e is not deletable. Then $(\rho, \varphi, \mathcal{T})$ is obviously feasible to $G - e$. Moreover $\tau^e(\rho, \varphi, \mathcal{T}) = \tau(\rho, \varphi, \mathcal{T})$. Indeed, by $d^\rho(e) = a(e) = 0$, we have $(d^\rho - a)^+(E \setminus e) = (d^\rho - a)^+(E)$. Also $d^\varphi(e) = 0$ implies $e \notin \delta X_\omega$ for $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$, which in turn implies $h(X_\omega) = h^e(X_\omega)$ by (3.24) (1). Thus $h^e \circ (\rho, \varphi, \mathcal{T}) = h \circ (\rho, \varphi, \mathcal{T})$.

Therefore $(\rho, \varphi, \mathcal{T})$ is feasible and not optimal to $G - e$. According to Lemmas 3.11 and 3.13, we can take a bipartite feasible triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ for $G - e$ such that

$$\tau^e(\rho^*, \varphi^*, \mathcal{T}^*) + 2 \leq \tau^e(\rho, \varphi, \mathcal{T}) = \tau(\rho, \varphi, \mathcal{T}) = \overline{\text{val}}(\mu, G, a) \text{ and,}$$

$$(d^{\rho^*} + d^{\varphi^*})(u, v) \leq (d^\rho + d^\varphi)(u, v) + 2 \quad (u, v \in V).$$

In particular,

$$d^{\rho^*}(e) + d^{\varphi^*}(e) \leq d^\rho(e) + d^\varphi(e) + 2 = 2.$$

So there are four cases $(d^{\rho^*}(e), d^{\varphi^*}(e)) = (0, 0), (2, 0), (1, 1), (0, 2)$. The first case is impossible. Otherwise $(\rho^*, \varphi^*, \mathcal{T}^*)$ is also feasible to G with $\tau(\rho^*, \varphi^*, \mathcal{T}^*) = \tau^e(\rho^*, \varphi^*, \mathcal{T}^*) < \overline{\text{val}}(\mu; G, a) \leq \tau(\rho^*, \varphi^*, \mathcal{T}^*)$; a contradiction.

Case 1: $(d^{\rho^*}(e), d^{\varphi^*}(e)) = (2, 0)$. In this case, triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ is also feasible to G (by $0 = d^{\varphi^*}(e) \leq |d^{\rho^*} - a|(e) = 2$). Hence

$$\tau^e(\rho^*, \varphi^*, \mathcal{T}^*) + 2 \leq \overline{\text{val}}(\mu; G, a) \leq \tau(\rho^*, \varphi^*, \mathcal{T}^*). \quad (3.27)$$

Moreover $(d^{\rho^*} - a)^+(E) = (d^{\rho^*} - a)^+(E \setminus e) + 2$, and $h \circ (\varphi^*, \mathcal{T}^*) = h^e \circ (\varphi^*, \mathcal{T}^*)$ by (3.24) (2) and $e \notin \delta X_\omega$ for $\omega \in \mathcal{E}_{\mathcal{T}^*, \varphi^*}$ (otherwise $d^{\varphi^*}(e) > 0$). Thus we have

$$\tau(\rho^*, \varphi^*, \mathcal{T}^*) = \tau^e(\rho^*, \varphi^*, \mathcal{T}^*) + 2. \quad (3.28)$$

By (3.27) and (3.28), $(\rho^*, \varphi^*, \mathcal{T}^*)$ is also optimal to G . In particular $(\rho^*, \varphi^*, \mathcal{T}^*)$ satisfies the optimality criterion (Lemma 3.12) with (ξ, f) . By (b) in Lemma 3.12, $0 = d^{\varphi^*}(e) < |d^{\rho^*} - a|(e) = 2$ implies $\xi(e) = 0$, which contradicts $\xi(e) > 0$.

Case 2: $(d^{\rho^*}(e), d^{\varphi^*}(e)) = (1, 1)$. Also in this case, triple $(\rho^*, \varphi^*, \mathcal{T}^*)$ is feasible to G . Hence (3.27) holds. Obviously $(d^{\rho^*} - a)^+(E) = (d^{\rho^*} - a)^+(E \setminus e) + 1$. There is unique $\omega \in \mathcal{E}_{\mathcal{T}^*, \varphi^*}$ with $\pi_\omega = 1$ and $e \in \delta X_\omega$. By (3.24), $h \circ (\varphi^*, \mathcal{T}^*) = h^e \circ (\varphi^*, \mathcal{T}^*) - h^e(X_\omega) + h(X_\omega)$. Hence we have

$$\tau(\rho^*, \varphi^*, \mathcal{T}^*) = \tau^e(\rho^*, \varphi^*, \mathcal{T}^*) + 1 + h^e(X_\omega) - h(X_\omega). \quad (3.29)$$

By (3.27) and (3.29), $h^e(X_\omega) = 1$ and $h(X_\omega) = 0$, and $(\rho^*, \varphi^*, \mathcal{T}^*)$ is also optimal to G . By $\pi_\omega > 0$, $h(X_\omega) = 0$, and (c) in Lemma 3.12, we have $\xi(\delta X_\omega \setminus \mathcal{D}^-(d^{\rho^*} - a)) + (1 - \xi)(\delta X_\omega \cap \mathcal{D}^-(d^{\rho^*} - a)) = 0$, implying

$$\xi(e') = \begin{cases} 0 & \text{if } e' \in \delta X_\omega \setminus \mathcal{D}^-(d^{\rho^*} - a), \\ 1 & \text{if } e' \in \delta X_\omega \cap \mathcal{D}^-(d^{\rho^*} - a). \end{cases}$$

Thus $\xi(e) = 0$ since $e \in \delta X_\omega \setminus \mathcal{D}^-(d^{\rho^*} - a)$. Again this contradicts $\xi(e) > 0$.

Case 3: $(d^{\rho^*}(e), d^{\varphi^*}(e)) = (0, 2)$. In this case, $(\rho^*, \varphi^*, \mathcal{T}^*)$ is not feasible to G . We modify $(\varphi^*, \mathcal{T}^*)$ as follows. Let $q := \varphi^*(x)$ and $q' := \varphi^*(y)$. Let $\bar{q} \in \mathcal{T}^*$ be the unique point with $d_{\mathcal{T}^*}(\bar{q}, q) = d_{\mathcal{T}^*}(q', \bar{q}) = 1$. We may assume $\bar{q} \prec q$. Also we may assume that \bar{q} is a vertex of \mathcal{T}^* ; attach a unit segment to \bar{q} if necessary. Then segments $\omega := (\bar{q}, q)$ and $\omega' := (\bar{q}, q')$ belong to $\mathcal{E}_{\mathcal{T}^*, \varphi^*}$. Delete open segment (\bar{q}, q) from \mathcal{T}^* , and identify q and q' . Let $(\tilde{\mathcal{T}}^*, \tilde{\varphi}^*)$ be the resulting metric-tree with a map. Then $(\rho^*, \tilde{\varphi}^*, \tilde{\mathcal{T}}^*)$ is bipartite feasible to G , since $d^{\tilde{\varphi}^*}(e) = 0$ and $d^{\tilde{\varphi}^*}(e') - d^{\varphi^*}(e') \in \{-2, 0\}$ ($e' \in E$). Hence we have

$$\tau^e(\rho^*, \varphi^*, \mathcal{T}^*) + 2 \leq \overline{\text{val}}(\mu; G, a) \leq \tau(\rho^*, \tilde{\varphi}^*, \tilde{\mathcal{T}}^*). \quad (3.30)$$

Obviously $(d^{\rho^*} - a)^+(E) = (d^{\rho^*} - a)^+(E \setminus e)$ (by $d^{\rho^*}(e) = 0 = a(e)$). Next we estimate $h^e \circ (\varphi^*, \mathcal{T}^*)$ and $h \circ (\tilde{\varphi}^*, \tilde{\mathcal{T}}^*)$. We first consider the case $\bar{q} \prec q'$. Let $X := X_\omega$ and $Y := X_{\omega'}$. Then $h \circ (\tilde{\varphi}^*, \tilde{\mathcal{T}}^*) = h^e \circ (\varphi^*, \mathcal{T}^*) - h^e(X) - h^e(Y) + h(X \cup Y)$ by construction. Hence we have

$$\tau(\rho^*, \tilde{\varphi}^*, \tilde{\mathcal{T}}^*) = \tau^e(\rho^*, \varphi^*, \mathcal{T}^*) + h^e(X) + h^e(Y) - h(X \cup Y). \quad (3.31)$$

By (3.30), $h^e(X) = h^e(Y) = 1$, $h(X \cup Y) = 0$, and $(\rho^*, \tilde{\varphi}^*, \tilde{\mathcal{T}}^*)$ is optimal to G . Consider edge $e' \in \delta X \setminus \{e\}$. If e' joins X and Y , then $d^{\tilde{\varphi}^*}(e') < d^{\varphi^*}(e') \leq |d^{\rho^*} - a|(e')$, and (b) in Lemma 3.12 is applicable. Otherwise $e' \in \delta(X \cup Y)$, and (c) in Lemma 3.12 is applicable with $h(X \cup Y) = 0$. Hence we have

$$\xi(e') = \begin{cases} 1 & \text{if } e' \in \mathcal{D}^-(d^{\rho^*} - a), \\ 0 & \text{otherwise,} \end{cases} \quad (e' \in \delta X \setminus \{e\}). \quad (3.32)$$

Decompose ξ into a convex combination of the incidence vectors of inner-odd-joins F_1, F_2, \dots, F_k . By (3.32), $F_i \cap (\delta X \setminus \{e\}) = \delta X \cap \mathcal{D}^-(d^{\rho^*} - a)$ for $i = 1, 2, \dots, k$. By $h^e(X) = 1$ and $e \in \delta X \setminus \mathcal{D}^-(d^{\rho^*} - a)$, we have $h(X) = 0$ ((3.24) (1)), implying that $|\delta X \setminus \mathcal{D}^-(d^{\rho^*} - a)|$ is even. Hence, for every $i = 1, 2, \dots, k$, we have

$$|\delta X| = |\delta X \cap \mathcal{D}^-(d^{\rho^*} - a)| + |\delta X \setminus \mathcal{D}^-(d^{\rho^*} - a)| \equiv |F_i \cap (\delta X \setminus \{e\})| \pmod{2}.$$

Here $|\delta X| \equiv |F \cap \delta X| \pmod{2}$ must hold for every inner-odd-join F and every $X \subseteq V \setminus S$. Hence $e \notin F_i$ for $i = 1, 2, \dots, k$, which implies $\xi(e) = 0$; a contradiction to the initial assumption. The argument for the case $q' \prec \bar{q}$ is essentially the same; replace $X \cup Y$ by $Y \setminus X$ in the argument above. The the proof of Proposition 3.10 is complete. \square

4 Polynomial solvability and NP-hardness

In this section, we prove the polynomial time solvability of μ -CEDP for a tree metric μ , and the NP-hardness of μ -EDP when μ is not a truncated tree metric.

4.1 Polynomial time solvability

Our approach is based on the polynomial equivalence between separation and optimization [12] as in [27]. We are given an instance $(G = (V, E), S, a, \mu)$ of μ -CEDP. Suppose that μ is a tree metric. Although the linear programming problem (2.2) has an exponential number of variables and inequalities, the multiflow-variable f has a compact edge-node formulation, and the separation of the inner-odd-join polytope Q can be done in polynomial time by Padberg-Rao's algorithm for minimum odd cuts [35]; see [28, Section 12.4] (and Exercise 8). Consequently the separation of (f, ξ) can be done in polynomial time. Thus, by the ellipsoid method [12], we can evaluate $\text{val}(\mu; G, a)$ in polynomial time. Consequently, for an edge $e \in E$, we can determine whether e is deletable or not. If e is deletable, then delete it. Repeat it at most $|E|$ times until all edges are not deletable. By Proposition 3.10, the resulting graph G is inner Eulerian. By any algorithm for the laminar locking theorem, e.g., [20], we can find an edge-disjoint set of S -paths attaining $\text{val}(\mu; G, 0)$ in polynomial time. This is our desired solution for μ -CEDP.

A generalization to the *capacitated* setting is straightforward. Now we are also given an integer-valued edge-capacity $c : E \rightarrow \mathbf{Z}_+$. The problem is to maximize $\sum_{P \in \Pi} f(P)(\mu(s_P, t_P) - a(P))$ over $f : \Pi \rightarrow \mathbf{Z}_+$ satisfying the capacity constraint $\sum_{P \in \Pi, e \in P} f(P) \leq c(e)$ for $e \in E$. Let $\text{val}(\mu; G, c, a)$ denote the maximum value. The LP-relaxation is constructed as follows. Let Q^c be the convex hull of all integer-valued functions $\xi : E \rightarrow \mathbf{Z}_+$ such that $0 \leq \xi(e) \leq c(e)$ for $e \in E$ and $\xi(\delta x) \equiv c(\delta x) \pmod{2}$ for $x \in V \setminus S$. In (2.2), replace $\xi(e) + \sum_{P \in \Pi, e \in P} f(P) \leq 1$ by $\xi(e) + \sum_{P \in \Pi, e \in P} f(P) \leq c(e)$, and replace $\xi \in Q$ by $\xi \in Q^c$. The resulting LP also has an integer optimal solution if μ is a tree metric. Since Q^c is a parity-constraint capacitated b -matching polytope, the separation for Q^c can also be done in polynomial time, as above. Thus we can evaluate $\text{val}(\mu; G, c, a)$ in polynomial time. Consequently, for each edge $e \in E$, we can determine the maximum deletion $\alpha^c(e) := \max\{\alpha \in [0, c(e)] \cap \mathbf{Z} \mid \text{val}(\mu; G, c - \alpha 1_e, a) = \text{val}(\mu; G, c, a)\}$ in polynomial

time by the bisection method, where 1_e is the incidence vector of e . The rest is the same as above.

4.2 NP-hardness

The goal of this subsection is to prove the NP-hardness of μ -EDP when μ is not a truncated tree metric (Theorem 1.3). We will prove two auxiliary results. The first one is a 4-point characterization of truncated tree metrics.

Theorem 4.1. *A weight μ is a truncated tree metric if and only if for all sets of four distinct elements $s, t, u, v \in S$, at least one of the following inequalities holds:*

$$\mu(s, t) = 0, \quad (4.1)$$

$$\mu(u, v) = 0, \quad (4.2)$$

$$\mu(s, t) + \mu(u, v) \leq \mu(s, v) + \mu(t, u), \quad (4.3)$$

$$\mu(s, t) + \mu(u, v) \leq \mu(s, u) + \mu(t, v). \quad (4.4)$$

Note that here we do not require the triangle inequality to hold. Therefore, if μ is not a truncated tree metric, then there are four elements s, t, u, v such that all inequalities (4.1)-(4.4) do not hold. From this violating quadruple (s, t, u, v) , we will reduce some NP-hard problem to $\mu|_{\{s, t, u, v\}}$ -EDP. For the reduction, we use the following version of the integer 2-commodity flow feasibility problem for which one of the commodities is unit:

Problem 4.2. The $(1, *)$ -edge-disjoint paths problem takes the input of an undirected graph $G = (V, E)$, nodes $r, q \in V$, and disjoint sets $R, Q \subseteq V$ such that $\{r\}, \{q\}, R, Q$ are disjoint and $|R| = |Q|$. The goal is to determine paths $P_0, P_1, \dots, P_{|R|}$ such that they are pairwise edge-disjoint, and P_0 connects r with q , and P_i connects a node in R with a node in Q , for $i = 1, \dots, |R|$, or establish that there is no such a family of paths.

Even, Itai, and Shamir [10] actually proved that a directed version of $(1, *)$ -edge-disjoint paths problem is NP-hard. Their reduction works for our undirected version with small change.

Theorem 4.3 (essentially [10]). *The $(1, *)$ -edge-disjoint paths problem is NP-hard.*

Now we prove Theorem 1.3 assuming Theorems 4.1 and 4.3, which will be proved later.

Proof of Theorem 1.3. Suppose that μ is not a truncated tree metric. Then Theorem 4.1 implies that there are four distinct elements s, t, u, v such that each of the following inequalities holds:

$$\mu(s, t) > 0, \quad (4.5)$$

$$\mu(u, v) > 0, \quad (4.6)$$

$$\mu(s, t) + \mu(u, v) > \mu(s, v) + \mu(t, u), \quad (4.7)$$

$$\mu(s, t) + \mu(u, v) > \mu(s, u) + \mu(t, v). \quad (4.8)$$

Then either both of

$$2\mu(s, t) + 2\mu(u, v) > \mu(s, u) + \mu(s, v) + \mu(u, v), \quad (4.9)$$

$$2\mu(s, t) + 2\mu(u, v) > \mu(t, u) + \mu(t, v) + \mu(u, v), \quad (4.10)$$

or both of

$$2\mu(s, t) + 2\mu(u, v) > \mu(s, t) + \mu(s, v) + \mu(t, v), \quad (4.11)$$

$$2\mu(s, t) + 2\mu(u, v) > \mu(s, t) + \mu(s, u) + \mu(t, u) \quad (4.12)$$

hold. Indeed, assume that at most one of (4.9)-(4.10), and at most one of (4.11)-(4.12) holds. By symmetry, we may assume that (4.9) and (4.11) do not hold. Summing them implies that

$$3\mu(s, t) + 3\mu(u, v) \leq \mu(s, u) + 2\mu(s, v) + \mu(t, v).$$

On the other hand, (4.5)-(4.8) imply that

$$\begin{aligned} 3\mu(s, t) + 3\mu(u, v) &> 2\mu(s, v) + 2\mu(t, u) + \mu(s, u) + \mu(t, v) \geq \\ &\geq \mu(s, u) + 2\mu(s, v) + \mu(t, v). \end{aligned}$$

This is a contradiction. So, by symmetry, we may assume that each of the inequalities (4.9)-(4.10) hold.

Now consider an instance of the $(1, *)$ -edge-disjoint paths problem, that is an undirected graph $G = (V, E)$, nodes $r, q \in V$, and disjoint sets $R, Q \subseteq V$, such that $\{r\}, \{q\}, R, Q$ are disjoint and $|R| = |Q|$. Define $G' := (V', E')$ by adding 4 new nodes to G , namely s, t, u, v , and add new edges ur and vq , and sx for all $x \in R$, ty for all $y \in Q$. A solution to the $\mu|_{S'}$ -EDP with $G', S' := \{s, t, u, v\}$ defines a set of paths so that each path has both its endpoints in S' . If this set of paths contains exactly 1 path from u to v , and $|R|$ paths from R to Q , then the weight is equal to $|R|\mu(s, t) + \mu(u, v)$. We claim that this routing – i.e. having 1 path from u to v , and $|R|$ paths from s to t – is the *only way* to achieve a weight of at least $|R|\mu(s, t) + \mu(u, v)$. This follows from the following claim.

Claim 4.4. Consider the perfect graph $K_{S'}$ with node set $S' := \{s, t, u, v\}$, and edge weights given by μ , and node-capacity $b(s) = b(t) = |R|$ and $b(u) = b(v) = 1$. Then $\xi(st) = |R|$ and $\xi(uv) = 1$ and $\xi(e) = 0$ for other edge e gives the unique maximum integral b -matching in $K_{S'}$.

Proof. Recall that an integral b -matching is a vector $\xi : E(K_{S'}) \rightarrow \mathbf{Z}_+$ such that $\xi \cdot \delta_{\{w\}} \leq b(w)$ for all $w \in S'$. Let ξ be a maximum weight integral b -matching with respect to weight $\mu|_{s,t,u,v}$. First suppose that $\xi(su) \geq 1$ and $\xi(tv) \geq 1$. Then, by (4.8), decreasing ξ on su and tv by one, and increasing x on st and uv by one will increase the weight of ξ . This is a contradiction. This implies that $\xi(su) = 0$ or $\xi(tv) = 0$. By symmetry we may assume $\xi(su) = 0$. Similarly we have (i) $\xi(sv) = 0$ or (ii) $\xi(ut) = 0$.

Case (i): $\xi(sv) = 0$. Now suppose that $\xi(tu) \geq 1$ and $\xi(tv) \geq 1$. Then decrease ξ on tu and tv by one, increase ξ on st by two, and increase ξ on uv by one. This, by (4.10), implies that the weight of ξ increases, a contradiction. Thus we get

that $\xi(tu) = 0$ or $\xi(tv) = 0$. Say, by symmetry, $\xi(tu) = 0$. To sum up, we now have $\xi(su) = \xi(sv) = x(tu) = 0$. If $\xi(tv) \geq 1$, then decrease ξ on tv by one, and increase ξ on st and uv by one, to experience the weight of ξ increased by (4.7). A contradiction, showing that $\xi(su) = \xi(sv) = \xi(tu) = \xi(tv) = 0$. By (4.5) and (4.6), and by the assumption that ξ has maximum weight, we get that $\xi(st) = |R|$ and $\xi(uv) = 1$.

Case (ii): $\xi(ut) = 0$. Since $b(v) = 1$, at most one of $\xi(sv)$ and $\xi(tv)$ is positive. By the same argument, we get $\xi(sv) = \xi(tv) = 0$, and hence $\xi(st) = |R|$ and $\xi(uv) = 1$, as required. This completes the proof. \square

Thus we conclude that there is a polynomial reduction from the $(1, *)$ -edge-disjoint paths problem to μ -EDP, implying that μ -EDP is NP-hard, proving Theorem 1.3. The remainder of this subsection is devoted to proving Theorems 4.1 and 4.3.

Characterization of truncated tree metrics. Here we prove Theorem 4.1. We start with a well-known 4-point characterization of tree metrics; see [43].

Theorem 4.5. *A weight μ is a tree metric if and only if it satisfies the triangle inequality, and for all sets of four distinct elements $s, t, u, v \in S$, at least one of the following inequalities holds:*

$$\mu(s, t) + \mu(u, v) \leq \mu(s, v) + \mu(t, u), \quad (4.13)$$

$$\mu(s, t) + \mu(u, v) \leq \mu(s, u) + \mu(t, v). \quad (4.14)$$

Hirai [14] considered the distance among *subtrees* in a tree, and derived a similar 4-point characterization for it. A weight μ is called a *subtree distance* if there are a metric-tree \mathcal{T} and a family $\{R_s\}_{s \in S}$ of its subtrees (connected subsets) indexed by S such that

$$\mu(s, t) = d_{\mathcal{T}}(R_s, R_t) \quad (s, t \in S).$$

In this case we say that $(\mathcal{T}, \{R_s\}_{s \in S})$ *realizes* μ or is a *realization* of μ . A truncated tree metric is exactly a subtree distance so that each subtree is a ball around some point. Hirai provided the following characterization of subtree distances.

Theorem 4.6 ([14]). *A weight μ is a subtree distance if and only if for all sets of four distinct elements $s, t, u, v \in S$, at least one of the following inequalities holds:*

$$\mu(s, t) = 0, \quad (4.15)$$

$$\mu(u, v) = 0, \quad (4.16)$$

$$\mu(s, t) + \mu(u, v) \leq \mu(s, v) + \mu(t, u), \quad (4.17)$$

$$\mu(s, t) + \mu(u, v) \leq \mu(s, u) + \mu(t, v), \quad (4.18)$$

$$2\mu(s, t) + 2\mu(u, v) \leq \mu(s, u) + \mu(s, v) + \mu(u, v), \quad (4.19)$$

$$2\mu(s, t) + 2\mu(u, v) \leq \mu(t, u) + \mu(t, v) + \mu(u, v), \quad (4.20)$$

$$2\mu(s, t) + 2\mu(u, v) \leq \mu(s, t) + \mu(s, v) + \mu(t, v), \quad (4.21)$$

$$2\mu(s, t) + 2\mu(u, v) \leq \mu(s, t) + \mu(s, u) + \mu(t, u). \quad (4.22)$$

Note that if a weight function satisfies (4.1)-(4.4), it will clearly also satisfy (4.15)-(4.22).

We are ready to prove Theorem 4.1. First we prove the “only if” part. Consider a realization $(\mathcal{T}, \{R_s\}_{s \in S})$ of the truncated tree metric as a subtree distance. Now each subtree R_s is a ball $B(p_s, r_s)$ with center p_s and radius r_s . Let $\bar{\mu}(s, t) := d_{\mathcal{T}}(p_s, p_t)$ for $s, t \in S$. Then $\bar{\mu}$ is a tree metric. Take any quadruple $s, t, u, v \in S$. We may assume that $\mu(s, t) > 0$ and $\mu(u, v) > 0$, for otherwise (4.1) or (4.2) holds. That implies $\mu(s, t) = \bar{\mu}(s, t) - r_s - r_t$ and $\mu(u, v) = \bar{\mu}(u, v) - r_u - r_v$. By Lemma 4.5, at least one of (4.13) and (4.14) holds for $\bar{\mu}$. Say, $\bar{\mu}(s, t) + \bar{\mu}(u, v) \leq \bar{\mu}(s, u) + \bar{\mu}(t, v)$, for example. Then $\mu(s, t) + \mu(u, v) = \bar{\mu}(s, t) + \bar{\mu}(u, v) - r_s - r_t - r_u - r_v \leq \bar{\mu}(s, u) + \bar{\mu}(t, v) - r_s - r_t - r_u - r_v \leq \mu(s, u) + \mu(t, v)$. This implies (4.4).

Second, using Lemma 4.6, we prove the “if” part. By Lemma 4.6, if μ satisfies at least one of (4.1)-(4.4) for every $s, t, u, v \in S$, then it must be a subtree distance. Now suppose that μ is realized by $(\mathcal{T}, \{R_s\}_{s \in S})$. Moreover we may assume that every leaf $p \in \mathcal{T}$ of the tree appears as a single-node subtree $R_s = \{p\}$ (for some $s \in S$) – otherwise we could chop off part of \mathcal{T} , and retain a realization of μ . Then we claim that every subtree R_s actually is a ball, which will prove the Lemma. For a fixed $s \in S$, if the diameter of R_s is zero, then we are done. Otherwise consider the median of R_s , that is a point $x \in R_s$ such that there are two distinct points x_1, x_2 in R_s at maximum distance from x . Let $r := d_{\mathcal{T}}(x, x_1) = d_{\mathcal{T}}(x, x_2) = \max_{x' \in R_s} d_{\mathcal{T}}(x, x')$. We can take leaves y_1, y_2 of \mathcal{T} such that $d_{\mathcal{T}}(y_1, y_2) = d_{\mathcal{T}}(y_1, R_s) + d_{\mathcal{T}}(x_1, x_2) + d_{\mathcal{T}}(R_s, y_2)$. We claim that $R_s = B(x, r)$. If by contradiction this were not true, then there would be a leaf y_3 such that $d_{\mathcal{T}}(R_s, y_3) > d_{\mathcal{T}}(B(x, r), y_3)$. Clearly $y_3 \neq y_1, y_2$. By our assumption, there must be s_1, s_2, s_3 such that $R_{s_1} = \{y_1\}$, $R_{s_2} = \{y_2\}$, $R_{s_3} = \{y_3\}$. Clearly, s, s_1, s_2, s_3 are four distinct elements of S . We claim that for $s = s, t = s_3, u = s_2, v = s_1$, none of the inequalities (4.1)-(4.4) holds: To check this, note that $y_1 \neq y_2$, which proves that (4.2) does not hold. From $d_{\mathcal{T}}(R_s, y_3) > d_{\mathcal{T}}(B(x, r), y_3)$ we get that $y_3 \notin R_s$, which proves that (4.1) does not hold. We also see that $\mu(s, t) + \mu(u, v) = d_{\mathcal{T}}(y_1, y_2) + d_{\mathcal{T}}(R_s, y_3) = d_{\mathcal{T}}(y_1, x) + d_{\mathcal{T}}(x, y_2) + d_{\mathcal{T}}(R_s, y_3) > d_{\mathcal{T}}(y_1, x) + d_{\mathcal{T}}(x, y_2) + d_{\mathcal{T}}(x, y_3) - r \geq d_{\mathcal{T}}(y_1, y_3) + d_{\mathcal{T}}(x, y_2) - r = d_{\mathcal{T}}(y_1, y_3) + d_{\mathcal{T}}(R_s, y_2) = \mu(s, u) + \mu(t, v)$, that is, (4.4) does not hold. A similar computation shows that (4.3) does not hold either. The contradiction proves that R_s is a ball for all $s \in S$, completing the proof of the Lemma. Based on the algorithm from Hirai [14], and the above proof, we get the following.

Corollary 4.7. *There is a polynomial time algorithm to decide whether a given weight μ is a truncated tree metric, and if so, also provide a realization $(\mathcal{T}, \{B(p_s, r_s)\}_{s \in S})$.*

Hardness of $(1, *)$ -edge-disjoint paths problems. The proof of Theorem 4.3 is a slight modification of [10]. Consider an instance of 3-SAT with variables x_1, \dots, x_n , and clauses c_1, \dots, c_m . Each clause c_j has three literals like x_i or \bar{x}_i .

Now we construct an instance of the $(1, *)$ -edge-disjoint paths problem. We introduce nodes r, q . For all $i = 1, \dots, n$, we introduce nodes s_i, t_i . For all $i = 1, \dots, n$ and $j = 1, \dots, m$, we introduce nodes $p_{2j-1}^i, p_{2j}^i, q_{2j-1}^i, q_{2j}^i$. We introduce edges to create paths $s_i, p_1^i, p_2^i, \dots, p_{2m}^i, t_i$, and edges to create paths $s_i, q_1^i, q_2^i, \dots, q_{2m}^i, t_i$. We add edges $rs_1, t_i s_{i+1}, t_n q$. For all $j = 1, \dots, m$, we

add nodes a^j, b_j , and for $k = 1, 2, 3$, we add nodes $b^{j,k}, a^{j,k}, a_{j,k}, b_{j,k}$ and edges $a^j b^{j,k}, b^{j,k} a^{j,k}, b_{j,k} a_{j,k}, a_{j,k} b_j$. For a literal $c_j = x_{u_1} \vee x_{u_2} \vee x_{u_3}$, and for $k = 1, 2, 3$, we add edges $a^{j,k} p_{2j-1}^{u_k}$ and $b_{j,k} p_{2j}^{u_k}$. If instead of x_{u_k} , we have $\overline{x_{u_k}}$ in literal c_j , then, instead, we add edges $a^{j,k} q_{2j-1}^{u_k}$ and $b_{j,k} q_{2j}^{u_k}$. Let $G = (V, E)$ denote the graph constructed this way. Let R' be the set of all nodes $a^j, a^{j,k}, a_{j,k}$, and let Q' be the set of all nodes $b_j, b^{j,k}, b_{j,k}$. Let R and Q be sets introducing a copy of a node in R' and Q' , respectively, and connect each of the nodes in R' and Q' with its copy.

The purpose of the construction is the following claim: the 3-SAT formula is satisfiable if and only if there is a set of edge-disjoint paths such that one of the paths goes from r to q , and the remaining paths are connecting a node in R with a node in Q . For the if part, consider a satisfying assignment. Then we use a path from s to t traversing all nodes s_i, t_i in this order, using the paths q_l^i if x_i was “true”, and using the paths p_l^i if x_i was “false”. For each j , clause c_j has a true literal $x_{u_k} = \text{“true”}$ or $\overline{x_{u_k}} = \text{“true”}$. Use path $a^{j,k}, p_{2j-1}^{u_k}, p_{2j}^{u_k}, b_{j,k}$ if $x_{u_k} = \text{“true”}$, and use path $a^{j,k}, q_{2j-1}^{u_k}, q_{2j}^{u_k}, b_{j,k}$ if $\overline{x_{u_k}} = \text{“true”}$. So we get a path connecting $a^{j,k}$ and $b_{j,k}$. Furthermore a^j is matched to $b^{j,k}$ and b_j is matched $a_{j,k}$ by one edge. Also for $k' \in \{1, 2, 3\} \setminus k$, $b^{j,k'}$ is matched to $a^{j,k'}$, and $b_{j,k'}$ is matched to $a_{j,k'}$. Finally add a perfect matching between R, Q and R', Q' . We get a solution of the instance of the $(1, *)$ -edge-disjoint paths problem. Conversely, suppose there is a solution to the $(1, *)$ -edge-disjoint paths problem. This contains one path P_0 from r to q , and a set of paths connecting each node in R with a node in Q . This path must go via the paths p_l^i or the paths q_l^i , for otherwise one of the terminals in R or Q would be cut off. We set $x_i = \text{“true”}$ if P_0 use q_l^i , and $x_i = \text{“false”}$ if P_0 use p_l^i . We show that this assignment is satisfiable. It is also straightforward that for each j , the copies of nodes $a^j, b_j, b^{j,k}, a^{j,k}, a_{j,k}, b_{j,k}$ ($k = 1, 2, 3$) are paired with each other, except for one of the nodes $a^{j,k}$, and one of the nodes $b_{j,k}$. In particular the unpaired node $a^{j,k}$ is necessarily incident to $p_{2j}^{u_k}$ or $q_{2j}^{u_k}$ not used by P_0 . This means that clause c_j has a true literal in this assignment for each j .

5 Concluding remarks

Here we discuss several related issues and raise open problems.

Node-disjoint version. As mentioned, Mader [34] extended (1.1) to a node-disjoint version. One may naturally ask a node-disjoint generalization of our results. Node-disjoint problems have a quite different nature from edge-disjoint problems. Indeed, an LP-based approach to Mader’s node-disjoint S -paths theorem is not known. Also there are algebraic generalizations to *group-labeled graphs* [8, 36].

In [25, Section 6], Karzanov announced a min-cost generalization of Mader’s node-disjoint S -paths theorem. This can be seen as a conjecture since his proof has not been completed (personal communication with A. Karzanov, 2009).

Recently Pap [38] proved a min-max theorem and the polynomial solvability of node-disjoint S -paths packing with truncated tree metric weights. His approach is quite different from that presented in this paper, and their relation is not clear. We do not know whether his approach can be extended to a min-cost version and whether it has a polyhedral interpretation.

Combinatorial polynomial time algorithm. Our algorithm crucially depends on the ellipsoid method. Karzanov's algorithm [23] for min-cost edge-disjoint S -paths is complicated, though it is combinatorial. So it is desirable to find a combinatorial polynomial time algorithm of a concise description. Such algorithms for disjoint S -paths packing are (i) (a reduction to) Lovasz's linear matroid matching [31, 42], (ii) Chudnovsky-Cunningham-Geelen's algorithm for non-zero A -paths [7], and (iii) Pap's algorithm for non-returning A -paths [37]. They solve node-disjoint problems (on group-labeled graphs for (ii) and (iii)). A natural approach is to modify or extend (i-iii) to our setting.

Here we suggest an alternative approach. As mentioned in the introduction, our formula includes a discrete minimax relation for inner-odd-joins F and tree locations ρ :

$$\max_F \min_{\rho} (d_{\Gamma, 1/2}^{\rho} - a)(E \setminus F) = \min_{\rho} \max_F (d_{\Gamma, 1/2}^{\rho} - a)(E \setminus F). \quad (5.1)$$

Here the LHS coincides with $\text{val}(\mu; G, a)$ by (2.1). So our problem also reduces to finding a *saddle point* in (5.1). Once a saddle point (F, ρ) is found, an optimal S -paths packing is obtained by solving an easy multiflow problem on an inner Eulerian graph $G - F$, according to Lemma 2.1. Such a game-theoretic consideration may bring a new algorithm to this problem.

Integrality gap of the LP-relaxation. The LP-relaxation (2.2) is applicable to μ -EDP/CEDP for any weight μ . Many existing approximation algorithms for μ -EDP use the natural LP-relaxation [4, 13, 45], which is known to have $\Omega(\sqrt{|V|})$ -integrality gap. Our LP-relaxation (2.2) is stronger than the natural LP. So it would be interesting to estimate the integrality gap of (2.2) and its dependency of μ , and to design approximation algorithms based on it. For example, consider the integer 2-commodity flow maximization $\mu_{2\text{com}}$ -EDP; see Introduction for definition of $\mu_{2\text{com}}$. By Rothschild-Winston theorem [40], for every inner Eulerian graph, the natural multiflow relaxation has an integral optimal solution with the value equal to the minimum 2-commodity cut. Thus the inequality (generally strict) holds:

$$\begin{aligned} \text{val}(\mu_{2\text{com}}; G, 0) &= \max_F \text{val}(\mu_{2\text{com}}; G - F, 0) = \max_F \min_X |\delta^{G-F} X| \\ &\leq \max_{\xi \in Q} \min_X (1 - \xi)(\delta^G X) \leq \min_X \max_F |\delta^{G-F} X|, \end{aligned}$$

where F denotes an inner-odd-join and X denotes a vertex subset with $s, s' \in X \not\ni t, t'$ or $s, t' \in X \not\ni t, s'$. Based on it, can we design a better approximation algorithm than the trivial 2-approximation algorithm?

Relation to Karzanov's approach. In [23], Karzanov established a min-max relation for μ -CEDP with $\mu = p1$ ($p > 0$); see also [25]. His min-max formula takes a form different from our min-max formula (1.2) or Theorem 3.5. Here we discuss their relationship. We need some notions to explain Karzanov's min-max formula. An *inner fragment* ϕ is a pair (X_{ϕ}, U_{ϕ}) of $X_{\phi} \subseteq V \setminus S$ and $U_{\phi} \subseteq \delta X_{\phi}$ with $|U_{\phi}|$ odd. Let \mathcal{F}_0 denote the set of all inner fragments. The characteristic

function χ_ϕ of an inner fragment ϕ is a function on E defined by

$$\chi_\phi(e) = \begin{cases} 1 & \text{if } e \in U_\phi, \\ -1 & \text{if } e \in \delta X_\phi \setminus U_\phi, \\ 0 & \text{otherwise,} \end{cases} \quad (e \in E).$$

Given $\beta : \mathcal{F}^0 \rightarrow \mathbf{R}_+$ and $\gamma : E \rightarrow \mathbf{R}_+$, define the function $l^{\beta,\gamma}$ on E by

$$l^{\beta,\gamma} := a + \gamma + \sum_{\phi \in \mathcal{F}^0} \beta(\phi) \chi_\phi.$$

We say that (β, γ) is *p-admissible* if $l^{\beta,\gamma} \geq 0$ and $l^{\beta,\gamma}(P) \geq p$ for all $P \in \Pi$.

Theorem 5.1 ([23]). *For any $p \geq 0$, we have*

$$\text{val}(p1; G, a) = \min \gamma(E) + \sum_{\phi \in \mathcal{F}^0} \beta_\phi(|U_\phi| - 1), \quad (5.2)$$

where the minimum is taken over all *p-admissible* pairs (β, γ) .

In fact, the minimization problem in RHS of (5.2) is the LP-dual of the LP obtained from (2.2) by substituting (2.6) to it and eliminating variable ξ . Then Theorem 5.1 follows from Theorem 3.9. Instead of detailed calculations, we explain how to obtain (β, γ) from our $(\rho, \varphi, \mathcal{T})$, and answer affirmatively Karzanov's question [25, Section 6, (2)]: *does there exist a half-integral optimal p-admissible pair (β, γ) when p and a are both integral?*

Suppose that p is even and a is even-valued. We show the existence of an integral optimal *p-admissible* pair (β, γ) . For each $s \in S$, consider a path P_s with length $p/2$ and ends q_s, p_s . Next identify q_s over all $s \in S$. The resulting tree (a subdivision of a star) is denoted by Γ . Then $(\Gamma, \{p_s\}_{s \in S}; 1)$ realizes $p1$. By Theorem 3.5, we can take an optimal bipartite feasible triple $(\rho, \varphi, \mathcal{T})$. Contract segment $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$ with $h[d^p - a](X_\omega) = 0$. Then $(\rho, \varphi, \mathcal{T})$ is an optimal (not necessarily bipartite) feasible triple such that d^p, d^φ , and π_ω are integral, and $h[d^p - a](X_\omega) = 1$ for each $\omega \in \mathcal{E}_{\mathcal{T}, \varphi}$.

Define (β, γ) by

$$\begin{aligned} \beta(\phi) &:= \begin{cases} \pi_\omega & \text{if } \exists \omega \in \mathcal{E}_{\mathcal{T}, \varphi} : X_\phi = X_\omega, U_\phi = \delta X_\omega \setminus \mathcal{D}^-(d^p - a), \\ 0 & \text{otherwise,} \end{cases} \quad (\phi \in \mathcal{F}^0), \\ \gamma(e) &:= \begin{cases} (d^p - a - d^\varphi)(e) & \text{if } e \notin \mathcal{D}^-(d^p - a), \\ 0 & \text{if } e \in \mathcal{D}^-(d^p - a), \end{cases} \quad (e \in E). \end{aligned}$$

Then $l^{\beta,\gamma}$ is given by

$$l^{\beta,\gamma}(e) = \begin{cases} d^p(e) & \text{if } e \notin \mathcal{D}^-(d^p - a), \\ (a - d^\varphi)(e) & \text{if } e \in \mathcal{D}^-(d^p - a), \end{cases} \quad (e \in E),$$

where we use $d^\varphi(e) = -\sum_\phi \beta(\phi) \chi_\phi(e)$ if $e \in \mathcal{D}^-(d^p - a)$ and $d^\varphi(e) = \sum_\phi \beta(\phi) \chi_\phi(e)$ if $e \notin \mathcal{D}^-(d^p - a)$. In particular $l^{\beta,\gamma}(e) \geq d^p(e) \geq 0$ (by $|d^p - a|(e) \geq d^\varphi(e)$). Hence

$l^{\beta, \gamma}(P) \geq d^{\rho}(P) = p$ for $P \in \Pi$, implying that (β, γ) is admissible. Moreover,

$$\begin{aligned} \gamma(E) + \sum_{\phi \in \mathcal{F}^0} \beta_{\phi}(|U_{\phi}| - 1) &= (d^{\rho} - a)^+(E) - d^{\varphi}(E \setminus \mathcal{D}^-(d^{\rho} - a)) \\ &\quad + \sum_{\omega \in \mathcal{E}_{\mathcal{T}, \varphi}} \pi_{\omega} \{ \delta_{X_{\omega}}(E \setminus \mathcal{D}^-(d^{\rho} - a)) - h[d^{\rho} - a](X_{\omega}) \} \\ &= (d^{\rho} - a)^+(E) - h[d^{\rho} - a] \circ (\varphi, \mathcal{T}) = \tau(\rho, \varphi, \mathcal{T}). \end{aligned}$$

Thus (β, γ) is optimal, and also integral by construction.

Acknowledgements

We thank Alexander Karzanov for remarks on the earlier version of the paper, and thank the referees for helpful comments. The first author is supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and is partially supported by Aihara Project, the FIRST program from JSPS. The second author is supported by the Hungarian National Foundation for Scientific Research (OTKA) grant CK80124.

References

- [1] Y. Bartal, On approximating arbitrary metrics by tree metrics, in: *Proceedings of the 30th Annual ACM Symposium on Theory of Computing (STOC 98)*, 161–168, ACM, New York, 1999.
- [2] M. Bădoiu, P. Indyk, and A. Sidiropoulos, Approximation algorithms for embedding general metrics into trees, in: *Proceedings of the Eighteenth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA 07)*, 512–521, ACM, New York, 2007.
- [3] L. Brunetta, M. Conforti, and M. Fischetti, A polyhedral approach to an integer multicommodity flow problem, *Discrete Applied Mathematics* **101** (2000), 13–36.
- [4] C. Chekuri, S. Khanna, and F.B. Shepherd, An $O(\sqrt{n})$ approximation and integrality gap for disjoint paths and unsplittable flow, *Theory of Computing* **2** (2006), 137–146.
- [5] B. V. Cherkassky, Solution of a problem of multiproduct flows in a network, *Ékonomika i Matematicheskie Metody* **13** (1977), 143–151, (in Russian).
- [6] V. Chepoi, F. Dragan, I. Newman, Y. Rabinovich, and Y. Vaxes, Constant approximation algorithms for embedding graph metrics into trees and outerplanar graphs, *Discrete & Computational Geometry* **47** (2012), 187–214.
- [7] M. Chudnovsky, W. H. Cunningham, and J. Geelen, An algorithm for packing non-zero A -paths in group-labelled graphs, *Combinatorica* **28** (2008), 145–161.
- [8] M. Chudnovsky, J. Geelen, B. Gerards, L. Goddyn, M. Lohman, and P. Seymour, Packing non-zero A -paths in group-labelled graphs, *Combinatorica* **26** (2006), 521–532.

-
- [9] J. Edmonds and E. L. Johnson, Matching, Euler tours and the Chinese postman, *Mathematical Programming* **5** (1973), 88–124.
- [10] S. Even, A. Itai, and A. Shamir, On the complexity of timetable and multicommodity flow problems, *SIAM Journal on Computing* **5** (1976), 691–703.
- [11] A. Frank, A. V. Karzanov, and A. Sebö, On integer multiflow maximization, *SIAM Journal on Discrete Mathematics* **10** (1997), 158–170.
- [12] M. Grötschel, L. Lovász, and A. Schrijver, *Geometric Algorithms and Combinatorial Optimization*, Springer-Verlag, Berlin, 1988.
- [13] V. Guruswami, S. Khanna, R. Rajaraman, B. Shepherd, and M. Yannakakis, Near-optimal hardness results and approximation algorithms for edge-disjoint paths and related problems, *Journal of Computer and System Sciences*, **67** (2003), 473–496.
- [14] H. Hirai, Characterization of the distance between subtrees of a tree by the associated tight span, *Annals of Combinatorics* **10** (2006), 111–128.
- [15] H. Hirai, Tight spans of distances and the dual fractionality of undirected multiflow problems, *Journal of Combinatorial Theory, Series B* **99** (2009), 843–868.
- [16] H. Hirai, Folder complexes and multiflow combinatorial dualities, *SIAM Journal on Discrete Mathematics* **25** (2011), 1119–1143.
- [17] H. Hirai, The maximum multiflow problems with bounded fractionality, RIMS-Preprint 1682, 2009.
- [18] H. Hirai, Half-integrality of node-capacitated multiflows and tree-shaped facility locations on trees, *Mathematical Programming, Series A*, to appear.
- [19] T. C. Hu, Multi-commodity network flows, *Operations Research* **11** (1963), 344–360.
- [20] T. Ibaraki, A. V. Karzanov, and H. Nagamochi, A fast algorithm for finding a maximum free multiflow in an inner Eulerian network and some generalizations, *Combinatorica* **18** (1998), 61–83.
- [21] A. V. Karzanov, Polyhedra related to undirected multicommodity flows, *Linear Algebra and its Applications* **114/115** (1989), 293–328.
- [22] A. V. Karzanov, Undirected multiflow problems and related topics—some recent developments and results, in: *Proceedings of the International Congress of Mathematician, Volume II*, Kyoto, Japan 1991, 1561–1571.
- [23] A. Karzanov, Edge-disjoint T -paths of minimum total cost, Report No. STAN-CS-92-1465, Department of Computer Science, Stanford University, Stanford, California, 1993, 66pp, available at <http://alexander-karzanov.net/>.
- [24] A. V. Karzanov, How to tidy up a symmetric set-system by use of uncrossing operations, *Theoretical Computer Sciences* **157** (1996), 215–225.
- [25] A. V. Karzanov, Multiflows and disjoint paths of minimum total cost, *Mathematical Programming* **78** (1997), 219–242.

- [26] A. V. Karzanov and M. V. Lomonosov, Systems of flows in undirected networks, in: *Mathematical Programming* (O.I. Larichev, ed.), Institute for System Studies, Moscow, 1978, 59–66 (in Russian).
- [27] J. C. M. Keijsper, R. A. Pendavingh, and L. A. Stougie, A linear programming formulation of Mader’s edge-disjoint paths problem, *Journal of Combinatorial Theory, Series B* **96** (2006), 159–163.
- [28] B. Korte and J. Vygen, *Combinatorial Optimization—Theory and Algorithms. Third Edition*, Springer-Verlag, Berlin, 2006.
- [29] L. Lovász, On some connectivity properties of Eulerian graphs, *Acta Mathematica Academiae Scientiarum Hungaricae* **28** (1976), 129–138.
- [30] L. Lovász, Selecting independent lines from a family of lines in a space, *Acta Scientiarum Mathematicarum (Szeged)* **42** (1980), 121–131.
- [31] L. Lovász, Matroid matching and some applications, *Journal of Combinatorial Theory, Series B* **28** (1980), 208–236.
- [32] N. Linial, E. London, and Y. Rabinovich, The geometry of graphs and some of its algorithmic applications, *Combinatorica* **15** (1995), 215–245.
- [33] W. Mader, Über die Maximalzahl kantendisjunkter A -Wege, *Archiv der Mathematik* **30** (1978), 325–336.
- [34] W. Mader, Über die Maximalzahl kreuzungsfreier H -Wege, *Archiv der Mathematik* **31** (1978/79), 387–402.
- [35] M. W. Padberg and M. R. Rao, Odd minimum cut-sets and b -matching, *Mathematics of Operations Research* **7** (1982), 67–80.
- [36] G. Pap, Packing non-returning A -paths, *Combinatorica* **27** (2007), 247–251.
- [37] G. Pap, Packing non-returning A -paths algorithmically, *Discrete Mathematics* **308** (2008), 1472–1488.
- [38] G. Pap, in preparation.
- [39] Y. Rabinovich and R. Raz, Lower bounds on the distortion of embedding finite metric spaces in graphs, *Discrete & Computational Geometry* **19** (1998), 79–94.
- [40] B. Rothschild and A. Whinston, On two-commodity network flows, *Operations Research* **14** (1966), 377–387.
- [41] A. Schrijver, A short proof of Mader’s \mathcal{S} -paths theorem, *Journal of Combinatorial Theory, Series B* **82** (2001), 319–321.
- [42] A. Schrijver, *Combinatorial Optimization—Polyhedra and Efficiency*, Springer-Verlag, Berlin, 2003.
- [43] C. Semple and M. Steel, *Phylogenetics*, Oxford University Press, Oxford, 2003.
- [44] P. D. Seymour, On odd cuts and plane multicommodity flows, *Proceedings of the London Mathematical Society* **42** (1981), 178–192.
- [45] V. V. Vazirani, *Approximation Algorithms*, Springer-Verlag, Berlin, 2001.